

SACCHARINITY

JAKOB KELLNER[†] AND SAHARON SHELAH[‡]

ABSTRACT. We present a method to iterate finitely splitting lim-sup tree forcings along non-wellfounded linear orders. We apply this method to construct a forcing (without using an inaccessible or amalgamation) that makes all definable sets of reals measurable with respect to a certain (non-ccc) ideal.

1. I

In the seminal paper [10] Solovay proved that in the Levy model (after collapsing an inaccessible) every definable set is measurable and has the Baire property.

In [9] Shelah showed that the inaccessible is necessary for measurability, but the Baire property of every definable set can be obtained by a forcing P without the use of an inaccessible (i.e. in ZFC). This forcing P is constructed by amalgamation of universally meager forcings Q . So every Q adds a co-meager set of generics and has many automorphisms, and the forcing P has similar properties to the Levy collapse. The property of Q that implies that Q can be amalgamated is called “sweetness” (a strong version of ccc).

One can ask about other ideals than Lebesgue-null and meager (or their defining forcings, random and Cohen), and classify such ideals (or forcings). This was done e.g. in Sweet & Sour [7] — sweet forcings are close to Cohen, sour ones close to random.

However, there is a completely different construction that brings measurability of all definable sets: Instead iterating basic forcings Q that have many automorphisms and add a measure 1 set of generics, we use a Q that adds only a null set of generics (a single one in our case, and this real remains the only generic over V_i even in the final limit). So Q has to be very non-homogeneous. Instead of having many automorphisms in Q , we assume that the skeleton of the iteration has many automorphisms (so in particular a non-wellfounded iterations has to be used).

We use the word Saccharinity for this concept: a construction that achieves the same effect as an (amalgamation of) sweet forcings, but using entirely different means.

As a first instance of this concept we construct an ideal I on the reals and a forcing P making every projective (or: definable) set measurable modulo I (without using amalgamation or an inaccessible). I will be defined from a tree forcing in the same way that the Marczewski ideal is defined from Sacks forcing. I is not a ccc ideal.

Annotated contents.

Section 2, p. 2: We define a class of tree forcings with “lim-sup norm”. The forcing conditions are subtrees of a basic tree that satisfy “along every branch, many nodes have many successors”.

Section 3, p. 6: We introduce a general construction to iterate such lim-sup tree-forcings along non-wellfounded total orders. It turns out that the limit is proper, ω^ω -bounding and has other nice properties similar to the properties of the lim-sup tree-forcings itself.

Date: November 2004.

1991 *Mathematics Subject Classification.* 03E35, 03E40.

[†] partially supported by FWF grant P17627-N12.

[‡] supported by the United States-Israel Binational Science Foundation (Grant no. 2002323), publication 859.

- Section 4, p. 14: We define (with respect to a lim-sup tree-forcing Q) the ideals $\mathbb{I}$ and $\mathbb{I}^c$ (the $< 2^{\aleph_0}$ -closure of $\mathbb{I}$). These ideals will generally not be ccc. We define what we mean by “ X is measurable with respect to $\mathbb{I}^c$ ” and formulate the main theorem: Assuming CH and a Ramsey property for Q (see Section 6), we can force that all definable sets are measurable.
- Section 5, p. 15: Assuming CH, we construct an order I which has many automorphisms and a certain cofinal sequence $(j_\alpha)_{\alpha \in \omega_2}$. We show that P (the nonwellfounded iteration of Q along the order I) forces that $2^{\aleph_0} = \aleph_2$, that $\mathbb{I}^c$ is nontrivial, that for every definable set X “locally” either all or no η_{j_δ} are in X and that the set $\{\eta_{j_\delta} : \delta \in \omega_2\}$ is of measure 1 in the set $\{\eta_i : i \in I\}$.
- Section 6, p. 20: We assume a certain Ramsey property for Q . We show that $\{\eta_i : i \in I\}$ is of measure 1 (with respect to $\mathbb{I}^c$). Together with the result of the last section this proves the main theorem.

2.

- -

We will define a class of tree forcings with “lim-sup norm”. Such trees (and generalizations) are investigated in [6] under the name $\mathbb{Q}_0^{\text{tree}}$, see Definition 1.3.5 there. Let us first introduce some notation around trees on $\omega^{<\omega}$.


Definition 2.1. Let $T \subseteq \omega^{<\omega}$ be a tree (i.e. T is closed under initial segments), let $s, t \in \omega^{<\omega}$, $A \subseteq T$.

- $\langle \rangle$ denotes the empty sequence.
- $s \leq t$ means that s is a restriction of t (or equivalently that $s \subseteq t$).
- t is immediate successor of s if $t \geq s$ and $\text{length}(t) = \text{length}(s) + 1$.
- $\text{succ}_T(t)$ is the set of immediate successors of t in a tree T . If the tree T is clear from the context we will also write $\text{succ}(t)$.
- s is compatible with t ($s \parallel t$), if $s \leq t$ or $t \leq s$. Otherwise, s and t are incompatible ($s \perp t$).
- $T^{[t]} := \{s \in T : s \parallel t\}$. (So $T^{[t]}$ is a tree.)
- $T \upharpoonright n := \{t \in T : \text{length}(t) < n\}$.
- $A \subseteq T$ is a chain if $s \parallel t$ for all $s, t \in A$.
- $b \subseteq T$ is a branch if it is a maximal chain.
- If there exists a $t \in b$ with $\text{length}(t) = n$ then this t is unique and denoted by $b \upharpoonright n$.
- $A \subseteq T$ is an antichain if $s \perp t$ for all $s \neq t \in A$.
- $A \subseteq T$ is a front if it is an antichain and every branch b meets A (i.e. $|b \cap A| = 1$).
- $t \leq A$ stands for: “ $t \leq s$ for some $s \in A$ ”.
- $T_{\text{cldn}}^A := \{t \in \omega^{<\omega} : t \leq A\}$.
- (We will use this notion only for finite antichains A . Then T_{cldn}^A is a finite tree.)
- If A and A' are antichains, then A' is stronger than A if for each $t \in A'$ there is a $s \in A$ such that $s \leq t$ (cf. Figure 1).
- If A and A' are antichains then A' is purely stronger than A if it is stronger and for each $s \in A$ there is a $t \in A'$ such that $s \leq t$ (cf. Figure 1).

We are only interested in finitely splitting trees (i.e. $\text{succ}(t)$ is finite for all $t \in T$). Then all fronts are finite. Note that being a front is stronger than being a maximal antichain. For example, $\{0^n 1 : n \in \omega\}$ is a maximal antichain in $2^{<\omega}$, but not a front.

Assumption. Assume T_{max} and μ satisfy the following:

- T_{max} is a finitely splitting tree.
- μ assigns a non-negative real to every subset of $\text{succ}_{T_{\text{max}}}(t)$ for every $t \in T_{\text{max}}$.
- μ is monotone, i.e. if $A \subseteq B$ then $\mu(A) \leq \mu(B)$.
- The measure of singletons is smaller than 1, i.e. $\mu(\{s\}) < 1$.
- For all branches b in T_{max} , $\limsup_{n \rightarrow \infty} (\mu(\text{succ}(b \upharpoonright n))) = \infty$.



- F 1. F' is stronger than F , F'' is purely stronger than F .

Such a $T_{\max}$ and μ define a lim-sup tree-forcing Q in the following way:

- Definition 2.2.**
- If T is a subtree of $T_{\max}$ and $t \in T$, then $\mu_T(t)$ is defined as the measure of the T -successors of t , i.e. $\mu_T(t) := \mu(\text{succ}_T(t))$.
 - Q consists of all subtrees T of $T_{\max}$ (ordered by inclusion) such that along every branch b in T , $\limsup(\mu_T(b \upharpoonright n)) = \infty$

So $T_{\max}$ itself is the weakest element of Q .

For example, Sacks forcing is such a forcing: Set $T_{\max} := 2^{<\omega}$, and for $t \in T_{\max}$ and $A \subseteq \text{succ}(t)$ set

$$\mu(A) := \begin{cases} \text{length}(t) & \text{if } |A| = 2, \\ 0 & \text{otherwise.} \end{cases}$$

Then a subtree T of $2^{<\omega}$ is in Q iff along every branch there are infinitely many splitting nodes.¹

Definition 2.3. A (finite or infinite) subtree T of $T_{\max}$ is n -dense if there is a front F in T such that $\mu_T(t) > n$ for every $t \in F$.

- Lemma 2.4.**
- (1) A subtree T of $T_{\max}$ is in Q iff T is n -dense for every $n \in \mathbb{N}$.
 - (2) “ $T \in Q$ ” and “ $T \leq_Q S$ ” are Borel statements, and “ $S \perp T$ ” is Π_1^1 (in the real parameters $T_{\max}$ and μ).

Proof. (1) $\rightarrow$: If $D_n := \{s \in T : \mu_T(s) > n\}$ meets every branch, then

$$F_n := \{s \in D : (\forall s' \not\leq s) s' \notin D\}$$

is a front.

$\leftarrow$: If b is a branch, then b meets every F_n , i.e. $\mu(b \upharpoonright m) > n$ for some m . Since $\mu(b \upharpoonright m)$ is finite, $\limsup(\mu_T(b \upharpoonright n))$ has to be infinite.

(2) Since $T_{\max}$ is finitely splitting, “ F is a front” is equivalent to “ F is a finite maximal antichain”. $\square$

A finite antichain A can be seen as an approximation of a tree: A approximates T means that A is a front in T . If A' is purely stronger than A , then A' gives more information about the tree T that is approximated (i.e. every tree approximated by A' is also approximated by A).

We will usually identify a finite antichain F and the corresponding finite tree T_{cln}^F .

- Definition 2.5.**
- A finite antichain F is n -dense if T_{cln}^F is n -dense.
 - $\bar{F} = (F_n)_{n \in \omega}$ is a front-sequence, if F_{n+1} is n -dense and purely stronger than F_n .
 - A front-sequence $\bar{F}$ and a tree $T \in Q$ correspond to each other if F_n is a front in T for all n .

¹This example is “discrete” in the following sense: For a node $s \in T$ there is an $A \subset \text{succ}(s)$ such that $\mu(A)$ is large but $\mu(B) < 1$ for every $B \subsetneq A$. In this paper, we will be interested in “finer” norms. In particular we will require the Ramsey property defined in 6.4.

- Facts 2.6.**
- If F is n -dense and F' is purely stronger than F , then F' is n -dense as well. (This is not true if F' is just stronger than F .)
 - If $T \in Q$ then there is a front-sequence corresponding to T .
 - If $\bar{F}$ is a front-sequence then there exists exactly one $T \in Q$ corresponding to $\bar{F}$, which we call $\lim(\bar{F})$. It is the tree

$$\lim(\bar{F}) := \{t \in T_{\max} : (\exists i \in \omega) t \leq F_i\},$$

or equivalently

$$\lim(\bar{F}) := \{t \in T_{\max} : (\forall i \in \omega) (\exists s \in F_i) t \parallel s\}.$$

Lemma 2.7. Assume that Q is a finitely splitting lim-sup tree-forcing.

- (1) If $T \in Q$ and $t \in T$ then $T^{[t]} \in Q$.
- (2) The finite union of elements of Q is in Q .²
- (3) The generic filter on Q is determined by a real η defined by $\Vdash_Q \{\eta\} = \bigcap_{T \in G_Q} \lim(T)$. It is forced that $\eta \notin V$ and that $T \in G_Q$ iff $\eta \in \lim(T)$. For every $T \in Q$ and $t \in T$ it is compatible with T that $t < \eta$ (i.e. $T \nVdash t \neq \eta$).
- (4) (Fusion) If $(T_i)_{i \in \omega}$ is a decreasing sequence in Q and $\bar{F}$ is a front-sequence such that F_i is a front in T_i for all i , then $\lim(\bar{F}) \leq_Q T_i$.
- (5) (Pure decision) If $D \subseteq Q$ is dense, $T \in Q$ and F is a front of T , then there is an $S \leq T$ such that F is a front of S and for every $t \in F$, $S^{[t]} \in D$.
- (6) Q is proper and ω^ω -bounding.

Sketch of proof. (1) and (2) and (4) are clear. (1) and (2) imply (5).

(3): Let G be Q -generic over V , and define $X := \bigcap_{T \in G} \lim(T)$. $\lim(T_{\max})$ is compact and therefore satisfies the finite intersection property. So X is nonempty. For every $T \in G$ and $n \in \omega$ there is exactly one $t \in T$ of length n such that $T^{[t]} \in G$. So X has at most one element.

If $r \in V$, then the set of trees $S \in Q^V$ such that $r \notin \lim(S)$ is dense: If r is a branch of $T \in Q$ then pick an m such that $\mu_T(r \restriction m) > 2$. Since singletons have measure less than 1, $r(m)$ has at least two immediate successors in T , and one of them (we call it t) is not an initial segment of r . So $S := T^{[t]}$ forces that $\eta \neq r$.

Assume towards a contradiction that $\eta \in \lim(T)$ for some $T \in Q^V \setminus G$. Then this is forced by some $S \in G$. In particular S can not be a subtree of T . So pick an $s \in S \setminus T$. Then $S^{[s]} \leq S$ forces that $\eta \notin \lim(T)$, a contradiction.

If $T \in Q$ and $t \in T$ then $T^{[t]}$ forces that $t < \eta$.

(4) and (5) imply that Q is ω^ω -bounding and satisfies a version of Axiom A (with fronts as indices instead of natural numbers).³ So we get properness. (We will prove a more general case in 3.25.) $\square$

So a front can be seen as a finite set of (pairwise incompatible) possibilities for initial segments of the generic real η . In the next section we will generalize this to sequences of generic reals instead of a single one. We will call sets of finite sequences of initial segments of these reals approximations.

Later we will need the following fact:


Lemma 2.8. If $S \in Q$ and the forcing R adds a new real $\mathfrak{r} \in 2^\omega$, then

² Q is generally not closed under countable unions.

³In the formulation of fusion and pure decision we could use the classical Axiom A version as well: Define F_n^T to be the minimal n -dense front, i.e.

$$F_n^T := \{t \in T : \mu_T(t) > n \text{ \& } (\forall s \preceq t) \mu_T(s) \leq n\},$$

and define $T \leq_n S$ by $T \leq S$ and $F_n^T = F_n^S$. It should be clear how to formulate fusion and pure decision for this notions, and that this proves Axiom A for Q . In 2.7 we implicitly use a different notion: $T \leq_A S$, meaning $T \leq S$ and A is a front in T and S . The reason is that this notion will be generalized for the non-wellfounded iteration.



F 2. An example for S and its subtree $\bar{T}$ (bold) when $r(0) = 0$.

R forces that there is a $T \leq_Q S$ such that $\lim(T) \cap V = \emptyset$, and moreover $\lim(T) \cap V$ remains empty in every extension of the universe.

Proof. Assume S corresponds to the front-sequence $\bar{F}$. Without loss of generality we can assume that along every branch in S there is exactly one split between F_{n-1} and F_n and this split has measure $> n$.

We define a name of a sequence of finite antichains (F'_n) the following way (cf. Figure 2): If n is even, we “take all splits”, i.e. F'_n is the set of nodes in F_n that are compatible with F'_{n-1} . If n is odd, then we add no splittings at all: for every $s \in F'_{n-1}$ we put exactly one successor $t \in F_n$ of s into F'_n , namely the one continuing the $r(\frac{n-1}{2})$ -th successor of the (unique) splitting node over s . It is clear that the sequence (F'_n) defines a subtree $\bar{T}$ of S .

Assume V' is an arbitrary extension of V containing an R -generic filter G over V . If $\eta \in \lim(\bar{T}[G]) \cap V$, then $r[G]$ can be decoded in V using S and η . This is a contradiction to $\Vdash_R r \notin V$. $\square$

We will also need the following family of definable dense subsets of Q :

Definition 2.9. Fix a recursive bijection ψ from ω to $2^{<\omega}$. Assume that $f : \omega \rightarrow \omega$ is strictly increasing and that $A \subseteq \omega$.

- For $g \in 2^\omega$, define $A_g^\psi := \{n \in \omega : \psi(n) < g\}$.
- Q_A^f is the set of all $T \in Q$ such that for all splitting nodes $t \in T$, $\text{length}(t)$ is in the interval $[f(n), f(n+1) - 1]$ for some $n \in A$.
- $T \in Q$ has full splitting with respect to f if for all $n \in \omega$ and $t \in T$ of length $f(n+1)$ there is an $s \leq t$ of length at least $f(n)$ such that $\mu_T(s) > n$.
- D_f^{spl} is the set of all $T \in Q$ such that either $T \in Q_{A_g^\psi}^f$ for some $g \in 2^\omega$ or $T \perp_Q S$ for all $g \in 2^\omega$ and $S \in Q_{A_g^\psi}^f$.

Of course the notions Q_A^f and D_f^{spl} depend on the forcing Q (i.e. on $T_{\max}$ and μ), so maybe it would be more exact to write $Q_A^f[T_{\max}, \mu]$ etc. However, we always assume that the Q is understood. The same applies to other notation in this paper (e.g. 4.1 or 4.3).

Lemma 2.10. Assume that $f : \omega \rightarrow \omega$ is strictly increasing and $A, B \subseteq \omega$.

- (1) If $g \neq g'$, then $A_g^\psi \cap A_{g'}^\psi$ is finite.
- (2) If A is finite then $Q_A^f = \emptyset$. $Q_\omega^f = Q$.
- (3) $Q_A^f \cap Q_B^f = Q_{A \cap B}^f$. If $A \subseteq B$, then $Q_A^f \subseteq Q_B^f$.
- (4) If $T \leq_Q S$ and $S \in Q_A^f$ then $T \in Q_A^f$.
- (5) For every $T \in Q$ there is a strictly increasing f such that T has full splitting with respect to f .

- (6) If $T \in Q$ has full splitting with respect to f and $|A| = \aleph_0$ then there is an $S \leq_Q T$ such that $S \in Q_A^f$.
- (7) D_f^{spl} is an (absolute definition of an) open dense subset of Q (using the parameters $f, T_{\max}$ and μ).⁴
- (8) In any extension V' of V the following holds: If $r \in 2^\omega \setminus V$ and $S \in Q_{A_r}^f$, then $T \perp_Q S$ for all $T \in V \cap D_f^{\text{spl}}$.

Proof. (1)–(4) and (6) are clear.

(5): Let T be an element of Q . Assume we already constructed $f(n)$. Let N be the maximum of $\mu_T(t)$ for $t \in T \restriction f(n)$. There is an $N + n + 1$ -dense front F in T . Let $f(n+1)$ be the maximum of $\{\text{length}(t) : t \in F\}$.

(7): “ T is incompatible with all $S \in Q_{A_g}^f$ ” is absolute, since it is equivalent to

$$(\forall g \in 2^\omega)(\forall S \subseteq T_{\max}) \left[S \notin Q_{A_g}^f \vee T \perp_Q S \right],$$

which is a Π_1^1 statement.

(8): Let $r \in 2^\omega \setminus V$ and $T \in V \cap D_f^{\text{spl}}$. If $T \in Q_{A_g}^f$ for some $g \in 2^\omega \cap V$, then $g \neq r$, so $A_g^\psi \cap A_r^\psi$ is finite and $Q_{A_r}^f \cap Q_{A_g}^f$ is empty. If on the other hand T is incompatible with all $S \in Q_{A_g}^f$ in V then this holds in V' as well. $\square$

Assume $f'(n) \geq f(n)$ for all $n \in \omega$. Define $h(n)$ by induction: $h(n+1) := f'(h(n)+1)$. If T has full splitting with respect to f , then T has full splitting with respect to h : $h(n) \leq f(h(n))$, since f is strictly increasing. $f(h(n)+1) \leq f'(h(n)+1) = h(n+1)$, and there are $h(n)$ -dense splits between the levels $f(h(n))$ and $f(h(n)+1)$. So there are n -dense splits between the levels $h(n)$ and $h(n+1)$. So we get:

Lemma 2.11. *If V' is an ω^ω -bounding extension of V and $T \in Q^{V'}$, then there is a strictly increasing $h \in V$ such that (in V') T has full splitting with respect to h .*

3. A - I

In this section we introduce a general construction to iterate lim-sup tree-forcings Q_i (as defined in the last section) along non-wellfounded linear orders I . It turns out that the limit P is proper, ω^ω -bounding and has other nice properties similar to the properties of Q_i itself. If I is wellfounded, then P is equivalent to the usual countable support iteration of (the evaluations of the definitions) Q_i .


There is quite some literature about non-wellfounded iteration. E.g. Jech and Groszek [3] investigated the wellfounded but non-linear iteration of Sacks forcings. Building on this,

⁴ $X_0 := \{A_g^\psi : g \in 2^\omega\}$ is an almost disjoint family, but not maximal. So of course $Q(X_0) := \bigcup_{A \in X_0} Q_A^f \subset Q$ is not dense. We add the incompatible conditions to get the dense set D_f^{spl} . One could ask whether $Q(X)$ is dense for a m.a.d. family X . The following holds:

(a) For every f there is a m.a.d. family X such that $Q(X)$ is not dense.
(b) (CH) For every f there is a m.a.d. family X such that $Q(X)$ is dense.
Proof: Fix f . A node $s \in T_{\max}$ has level m if $f(m) \leq \text{length}(s) < f(m+1)$. $S \in Q$ has unique splitting if S has at most one splitting point of level n for all $n \in \omega$. For every $T \in Q$ there is an $S \leq_Q T$ with unique splitting.

For (a), fix a $T \in Q$ with unique splitting. Set $Y := \{A \in [\omega]^{\aleph_0} : (\forall S \leq_Q T) S \notin Q_A^f\}$. Y is open dense in $([\omega]^{\aleph_0}, \subseteq)$, therefore there is a m.a.d. $X \subseteq Y$.

For (b), list Q as $(T_\alpha)_{\alpha \in \omega_1}$, and build $B_\alpha \in [\omega]^{\aleph_0}$ by induction on $\alpha \in \omega_1$: Find an $S \leq_Q T_\alpha$ with unique splitting. If some $S' \leq_Q S$ is in $Q_{B_\beta}^f$ ($\beta < \alpha$) (or equivalently in $Q_{\bigcup_{i \in I} B_{\beta_i}}^f$ for some $I \in \omega, \beta_0, \dots, \beta_{I-1} < \alpha$), then just pick any almost disjoint B_α . Otherwise enumerate $(B_\beta)_{\beta \in \alpha}$ as $(C_n)_{n \in \omega}$, and construct B_α and $S' \leq_Q S$ inductively: At stage n , add a split of S to S' whose level is not in $\bigcup_{m \leq n} C_m$, and use some bookkeeping to guarantee that $S' \in Q$. Let B_α be the set of splitting-levels of S' .



F 3. An approximation g ; $u = \{i, j\}$, $T_{\max}^i = 2^{<\omega}$, $T_{\max}^j = 3^{<\omega}$.
 $\text{Pos}(g) = \text{Pos}_{\leq j}(g) = \{(a_i^1, b_j^0), (a_i^1, b_j^1), (a_i^0, b_j^2), (a_i^0, b_j^3), (a_i^0, b_j^4)\}$.

Kanovei [5] and Groszek [2] develop non-wellfounded iterations of Sacks. The description of the construction there is very close to what we do here, but the realization is quite different. Brendle [1] presented finite-support non-wellfounded iteration constructions (extracted from Shelah's method of iterations along smooth templates [8]).

Definition 3.1. Let I^∞ be a linear order containing a maximal element ∞ . For $i \in I^\infty$ we set $I_{<i} := \{j \in I : j < i\}$ and $I := I_{<\infty}$.

For every $i \in I$ we fix a finitely splitting lim-sup tree-forcing Q_i (to be more exact, fix a pair $(T_{\max}^i, \mu^i)$). In the application of this paper, each Q_i will be the same forcing Q .

Definition 3.2. (Pre-condition) We call p a pre-condition, if p is a function, the domain of p is a countable⁵ subset of I , and for each $i \in \text{dom}(p)$, $p(i)$ consists of the following:

- Dom_i^p , a countable subset of $\text{dom}(p) \cap I_{<i}$, and
- a (definition of a) Borel function $B_i^p : (\omega^\omega)^{\text{Dom}_i^p} \rightarrow Q_i$

The idea is that we calculate the condition $B_i^p \in Q_i$ using countably many generic reals $(\eta_j)_{j \in \text{Dom}_i^p}$ that have already been produced at stage i . We will only be interested in B 's that are continuous (on a certain Borel set), i.e. if we want to know B_i^p up to some finite height we only have to know $(\eta_i \upharpoonright m)_{i \in u}$ for some finite u and $m \in \omega$. Moreover, we will assume that we will have “wellfounded continuity parts”:

Definition 3.3. (Approximation)

- g is an approximation, if g is a function with finite domain $u \subseteq I$ of the following form: Let i_0 be the smallest element of u . $\text{Pos}_{<i_0}(g) := \{\emptyset\}$, and (by induction on $i \in u$): $g(i)$ is a function from $\text{Pos}_{<i}(g)$ to finite antichains in $T_{\max}^i$, and $\text{Pos}_{\leq i}(g) := \{\bar{a} \cup (i, a_i) : \bar{a} \in \text{Pos}_{<i}(g), a_i \in g(i)(\bar{a})\}$.
- For any $i \in I^\infty$, $\text{Pos}_{<i}(g)$ is defined as $\text{Pos}_{\leq j}(g)$, where $j = \max(\text{dom}(g) \cap I_{<i})$ (or as $\{\emptyset\}$, if $\text{dom}(g) \cap I_{<i}$ is empty). $\text{Pos}(g) := \text{Pos}_{<\infty}(g)$. $\text{Pos}(g)$ is called the set of possibilities of g .
- If $i \notin \text{dom}(g)$ or $\bar{a} \notin \text{Pos}_{<i}(g)$ we set $g(i)(\bar{a}) := \{\langle \rangle\}$ (i.e. the front in $T_{\max}$ consisting only of the root. This corresponds to “no information”).
- Let g be an approximation, $J \subset I$, and $\bar{\eta} = (\eta_i)_{i \in J}$ a sequence of reals. Then $\bar{\eta}$ is compatible with g , if there is an $\bar{a} \in \text{Pos}(g)$ such that $a_i < \eta_i$ for all $i \in \text{dom}(g) \cap J$. If in addition $J \supseteq \text{dom}(g)$, then this $\bar{a}$ is uniquely defined, and we set $g(i)(\bar{\eta}) := g(i)(\bar{a} \upharpoonright I_{<i})$.

In the same way we define $\bar{b}$ to be compatible with g for $\bar{b} = (b_i)_{i \in J}$ a sequence of elements of $\omega^{<\omega}$. Then $g(i)(\bar{b}) := g(i)(\bar{a} \upharpoonright I_{<i})$ is well-defined if $J \supseteq \text{dom}(g)$ and the elements of $\bar{b}$ have sufficient length.

⁵this includes finite and empty.

- If g and g' are both approximations, then g' is stronger than g if $\text{dom}(g') \supseteq \text{dom}(g)$ and for all $\bar{b} \in \text{Pos}(g')$ there is an $\bar{a} \in \text{Pos}(g)$ such that $\bar{b} \geq \bar{a}$ (i.e. $b_i \geq a_i$ for all $i \in \text{dom}(g)$). In this case $\bar{a}$ is uniquely defined and called $\bar{b} \upharpoonright g$.
- g' is purely stronger than g if g' is stronger than g and for all $i \in \text{dom}(g)$ and $b \in \text{Pos}_{<i}(g')$ the front $g'(i)(\bar{b})$ is purely stronger than $g(i)(\bar{b} \upharpoonright g)$.
- For $u \subseteq \text{dom}(g)$, $\text{maxlength}_u(g)$ is $\max(\{\text{length}(a_i) : i \in u, \bar{a} \in \text{Pos}(g)\})$. $\text{maxlength}(g)$ is $\text{maxlength}_{\text{dom}(g)}(g)$. $\text{minlength}(g)$ is defined analogously.
- g is n -dense for $i \in I$, if $i \in \text{dom}(g)$ and for all $\bar{a} \in \text{Pos}_{<i}(g)$, $g(i)(\bar{a})$ is n -dense for Q_i (see 2.5).
- For all $\bar{a} = (a_i)_{i \in u}$ such that $a_i \in T_{\max}^i$ there is a (unique) approximation $\bar{g}$ such that $\text{Pos}(g) = \{\bar{a}\}$. We will call this approximation $\bar{a}$ as well.

So if g' is stronger than g , then all $\bar{\eta}$ compatible with g' are compatible with g . Clearly “stronger” and “purely stronger” are transitive.

Definition 3.4. (Approximation to p) Let p be a pre-condition.

- g approximates p , or: g is a p -approximation, if $\text{dom}(g) \subseteq \text{dom}(p)$ and g is an approximation with the following property: If $i \in \text{dom}(g)$, $\bar{a} \in \text{Pos}_{<i}(g)$, and $(\eta_j)_{j \in \text{Dom}_i^p}$ is compatible with $\bar{a}$, then $g(i)(\bar{a})$ is a front in $B_i^p(\bar{\eta})$.
- g is an indirect approximation to p witnessed by g' , if g' approximates p and g' is purely stronger than g .

Fact 3.5. If g (indirectly) approximates p , g' approximates p , and g' is stronger than g , then g' is purely stronger than g .

Now we can define the forcing P , the non-wellfounded countable support limit along I :

Definition 3.6. (The nwf-iteration $P = \text{nwf-lim}_I(Q_i)$)

- $p \in P$ means:
 p is a pre-condition, and for all $u \subseteq \text{dom}(p)$, $i \in u$ and $n \in \omega$ there is a p -approximation g such that $\text{dom}(g) \supseteq u$, g is n -dense for i , and $\text{minlength}_u(g) > n$.
- For $p, q \in P$, $q \leq p$ means:
for all p -approximations g there is a q -approximation h which is stronger than g (so in particular, $\text{dom}(q) \supseteq \text{dom}(p)$).
- $q \leq_g p$ if $q \leq p$ and g indirectly approximates p and q .

As we will see in 3.25(6), P is just the usual countable support iteration of Q_i in case that I is well-ordered.

- Facts 3.7.**
- $\leq$ is transitive, and for a fixed approximation g the relation $\leq_g$ is transitive as well. Also, if g' is purely stronger than g then $\leq_{g'}$ implies $\leq_g$.
 - For every $p \in P$, the approximations of p are directed: If g and g' both (indirectly) approximate p , then there is a g'' approximating p that is (purely) stronger than both g and g' (and every p -approximation g'' has this property if $\text{dom}(g'') \supseteq \text{dom}(g) \cup \text{dom}(g')$ and if $\text{minlength}_{\text{dom}(g) \cup \text{dom}(g')}(g'')$ is large enough).

So in particular for every $p \in P$ there is an approximating sequence:

Definition 3.8. An approximating sequence for $p \in P$ is a sequence $(g_n)_{n \in \omega}$ of approximations of p such that g_{n+1} is purely stronger than g_n , g_{n+1} is n -dense for each $i \in \text{dom}(g_n)$, and $\text{dom}(p) = \bigcup \text{dom}(g_n)$.

An approximating sequence contains all relevant information about p . In particular, g is an indirect approximation to p iff there is an n such that g_n is purely stronger than g . So if p and q both have the approximating sequence $(g_n)_{n \in \omega}$, then $p \leq q$ and $q \leq p$ (and $p \leq_g q$ if g indirectly approximates p or q).

Approximating sequences provide an equivalent definition for P :

Definition 3.9. (Alternative definition of the nwf-iteration P) Define the p.o. P' as follows: $\bar{g} \in P'$ iff $\bar{g}$ is a sequence of approximations $(g_n)_{n \in \omega}$ such that g_{n+1} is purely stronger than g_n and n -dense for every $i \in \text{dom}(g_n)$.

$\bar{h} < \bar{g}$ iff for every n there is an m such that h_m is stronger than g_n .

Lemma 3.10. *There is a dense embedding⁶ $\phi : P' \rightarrow P$.*

Proof. Given a sequence $\bar{g} \in P'$, define $p = \phi(\bar{g})$ the following way: $\text{dom}(p) = \bigcup \text{dom}(g_n)$. For $i \in \text{dom}(p)$, set $\text{Dom}_i^p := \text{dom}(p) \cap I_{<i}$. Define $T = B_i^p(\bar{\eta})$ as follows: If $\bar{\eta}$ is compatible with all g_n , then let T be $\{t \in T_{\max}^i : (\exists n \in \omega) t \leq g_n(i)(\bar{\eta})\}$. Otherwise, let n be maximal such that $\bar{\eta}$ is compatible with g_n , and let T be $\{t \in T_{\max}^i : (\exists s \in g_n(i)(\bar{\eta})) t \parallel s\}$. Clearly, $B_i^p \in Q_i$, B_i^p is a Borel function, and each g_n approximates p . Therefore $(g_n)_{n \in \omega}$ is an approximating sequence for $p \in P$. It is clear that ϕ preserves the order.

Let ψ map $p \in P$ to any approximating sequence for p . $\psi : P \rightarrow P'$ preserves order as well and $\phi(\psi(p)) =^* p$. Therefore ϕ is a dense embedding. $\square$

To summarize: Every $p \in P$ corresponds to a purely increasing sequence (g_n) of approximations such that $\bigcup \text{dom}(g_n) = \text{dom}(p)$ and g_{n+1} is n -dense for $\text{dom}(g_n)$. The approximating sequences immediately prove a version of fusion:

Lemma 3.11. (Fusion) *Assume that $(p_n)_{n \in \omega}$ is a sequence of conditions, $(g_n)_{n \in \omega}$ a sequence of approximations, and $i_n \in \text{dom}(g_n)$ such that:*

- $p_{n+1} \leq_{g_n} p_n$,
- g_{n+1} is purely stronger than g_n and n -dense for i_n ,
- $(i_n)_{n \in \omega}$ covers $\bigcup \text{dom}(p_n)$ infinitely often.

Then there is a condition p_ω such that $p_\omega \leq_{g_n} p_n$ for all n .

Proof. We already know that the sequence $(g_n)_{n \in \omega}$ of approximations defines a condition p_ω such that each g_n approximates p_ω . If h approximates p_n , then some g_m is stronger than h . g_m approximates p_ω , so $p_\omega \leq p_n$. $\square$


- Notes 3.12.**
- (1) If g indirectly approximates p , then there is a $q =^* p$ such that g approximates q . (Just let q correspond to an approximating sequence of p starting with $g_0 = g$.)
 - (2) It doesn't matter whether the g_n in an approximating sequence are approximations to p or just indirect approximations.
 - (3) It doesn't matter whether g_{n+1} proves n -density for every $i \in \text{dom}(g_n)$ or for just some i_n , provided that the sequence $(i_n)_{n \in \omega}$ covers $\bigcup \text{dom}(g_n)$ infinitely often.
 - (4) In Definition 3.2 of pre-condition, instead of requiring B_i^p to be a function into Q , we could have defined B_i^p to be a function to subtrees of $T_{\max}$. The additional “ n -dense” requirements on a condition guarantee $B_i^p(\bar{\eta}) \in Q$ anyway (for generic sequences $\bar{\eta}$).
 - (5) Every approximation g can be interpreted as a condition in P , by

$$B_i^g(\bar{\eta}) := \{t : t \parallel g(i)(\bar{\eta})\} \text{ for } i \in \text{dom}(g).$$

(Recall that $g(i)(\bar{\eta}) := \{\langle \rangle\}$ if $\bar{\eta}$ is incompatible with g .) Then g approximates itself.

- (6) For any approximation g and $u \subseteq I$ finite we can assume $u \subseteq \text{dom } g$: Just set $g(i) = \{\langle \rangle\}$ for $i \notin \text{dom } g$. This is consistent with the notational convention in 3.3.
- (7) If g and h are approximations, we can assume without loss of generality that $\text{dom}(g) = \text{dom}(h)$.
- (8) For any $U \subseteq I$ countable and $p \in P$ we can assume without loss of generality that $\text{dom}(p) \supseteq U$. This is clear if p is interpreted as a sequence of Borel-functions (just set $B_i^p = T_{\max}^i$ for $i \notin \text{dom}(p)$). If p is interpreted as sequence $(g_n)_{n \in \omega}$ of approximations, we have to set $g_n(i)$ to be $T_{\max}^i \restriction k(n)$ for some sufficiently large $k(n)$. (Using $\{\langle \rangle\}$ does not work here, since it does not prove n -density.)

⁶ ϕ is even an isomorphism modulo $=^*$, where $p =^* q$ if $q \leq p$ and $q \leq p$.



F 4.

We now investigate conjunctions of antichains (cf. Figure 4), of approximations and of a condition and an approximation.

Definition 3.13. • Let F and F' be finite antichains in $T_{\max}^i$. Set

$$A := \{t \in F : (\exists s \in F') t \geq s\} \cup \{s \in F' : (\exists t \in F) s \geq t\}.$$

Either A is empty (then F and F' are called incompatible, and $F \wedge F'$ is undefined), or A is a nonempty antichain (then F and F' are compatible, and $F \wedge F' := A$).

- Now we define $h = g \wedge g'$ for two approximations g and g' . Without loss of generality⁷ $\text{dom}(g) = \text{dom}(g') = u$. Let h be the approximation with domain u defined inductively for all $i \in u$: For $\bar{t} \in \text{Pos}_{<i}(h)$ set

$$h(i)(\bar{t}) := g(i)(\bar{t} \upharpoonright \text{dom}(g)) \wedge g'(i)(\bar{t} \upharpoonright \text{dom}(g')), \text{ if defined.}$$

If at any $i \in u$ this $h(i)$ is undefined, then g, g' are called incompatible and $g \wedge g'$ is undefined. Otherwise set $g \wedge g' := h$.

Facts 3.14. • If there is an approximation stronger than both h and g , then $h \wedge g$ is defined; and if $h \wedge g$ is defined then it is the weakest approximation stronger than both h and g .

- If g approximates p , h approximates q , and h, g are incompatible, then $p \perp_Q q$.
- If g' is stronger than g , then $h \wedge g'$ (if defined) is stronger than $h \wedge g$.

Notes. • It is possible that $h \wedge g$ is defined but $h \wedge g'$ is not for some g' stronger than g .

- If g' is purely stronger than g , and $h \wedge g'$ is defined, then $h \wedge g'$ does *not* have to be purely stronger than $h \wedge g$ (see Figure 4).
- However, if for $u := \text{dom}(h) \cap \text{dom}(g)$, $\text{minlength}_u(g) > \text{maxlength}_u(h)$, if $g \wedge h$ exists and if g' is purely stronger than g , then $g' \wedge h$ exists and is purely stronger than $g \wedge h$.

It is possible to define $q \wedge p$ for $p \parallel q \in P$ (using some kind of Cantor Bendixon algorithm). However, we will only use the version $p \wedge h$ for an approximation h : Assume that p corresponds to the sequence (g_n) of approximations. If for some n , h and g_n are incompatible, then there is no $q \leq p, h$. On the other hand, if h is compatible with all g_n , then for n large enough $g_{n+1} \wedge h$ is purely stronger than $g_n \wedge h$ and n -dense for all $i \in \text{dom}(g_n)$. So $(g_n \wedge h)_{n \in \omega}$ is an approximating sequence corresponding to some $p \wedge h \in P$.

Definition 3.15. Let p correspond to the approximating sequence $\bar{g}$. p is compatible with h if h and g_n are compatible for all n . In this case $p \wedge h \in P$ is defined by the approximating sequence $(g_n \wedge h)_{n \geq m}$ for some suitable m .

Then we get the following:

Facts 3.16. Let $p \in P$ be compatible to the approximation h .

⁷According to 3.12(7).

- $p \wedge h \in P$, $p \wedge h \leq p$. If h (indirectly) approximates p , then $p \wedge h =^* p$.
- If g indirectly approximates p and the domain and length of g is large enough, then $g \wedge h$ indirectly approximates $p \wedge h$.
- If $q \leq p$ and g is compatible to q , then g is compatible to p and $q \wedge g \leq p \wedge g$.
- If h is stronger than g and h is compatible to p , then g is compatible to p and $p \wedge h \leq p \wedge g$.
- If $q \leq p$ and g indirectly approximates q then g is compatible with p and $q \leq p \wedge g$.

An important special case is that of a sub-approximation:

Definition 3.17. h is sub-approximation of g if $\text{Pos}(h) \subseteq \text{Pos}(g)$.

In particular a sub-approximation of g is stronger than g .

Lemma 3.18. (Sub-approximation) Assume that g indirectly approximates p and that h is a sub-approximation of g . Then h is compatible with p . If $q \leq p$ and h approximates q , then there is an $r \leq_g p$ such that $r \wedge h =^* q$.

Proof. We will define r so that it is identical to q “below h ” and identical to p otherwise.

More exactly: Without loss of generality we can assume that p corresponds to $(g_n)_{n \in \omega}$, $g_0 = g$, q corresponds to $(h_n)_{n \in \omega}$, $h_0 = h$, h_n is stronger than g_n , and $\text{dom}(h_n) = \text{dom}(g_n) = u_n$. Let f_n be an approximation with domain u_n , i_0 minimal in u_n . So $\text{Pos}_{< i_0}(f_n) = \{\emptyset\}$. By induction on $i \in u_n$, define for all $\bar{a} \in \text{Pos}_{< i}(f_n)$

$$f_n(i)(\bar{a}) := \begin{cases} g_n(i)(\bar{a} \upharpoonright g_n) & \text{if } \bar{a} \text{ is incompatible with } h_n, \\ h_n(i)(\bar{a}) \cup \{t \in g_n(i)(\bar{a} \upharpoonright g_n) : t \perp h_n(i)(\bar{a})\} & \text{otherwise.} \end{cases}$$

It is clear that the possibilities of f_n follow h_n up to some $i \in \text{dom } g_n$ and from then on follow g_n . To be more exact: $\bar{a} \in \text{Pos}(f_n)$ iff $\bar{a} \upharpoonright g_n \in \text{Pos}(g_n)$ and for some $i \in \text{dom}(g_n) \cup \{\infty\}$, $\bar{a} \upharpoonright I_{< i}$ is in $\text{Pos}(h_n)$ and either $i = \infty$ or $a_i \perp h_n(i)(\bar{a})$. From this it follows that f_n is purely stronger than g_n , and that the f_n are an approximating sequence (converging to some $r \leq p$). So $r \leq_g p$, $f_n \wedge h = h_n$, and $r \wedge h =^* q$. $\square$

A special case of a sub-approximation is a singleton:

Definition 3.19. Assume that g indirectly approximates p and $\bar{a} \in \text{Pos}(g)$. Then the approximation $\bar{a}$ is compatible with p . Define $p^{[\bar{a}]}$ to be $p \wedge \bar{a}$.

So we get:

Corollary 3.20. (Specialization and pure decision) Assume that g indirectly approximates p and that $\bar{a} \in \text{Pos}(g)$.

- (1) $p^{[\bar{a}]} \in P$, $p^{[\bar{a}]} \leq p$ and $\bar{a}$ indirectly approximates $p^{[\bar{a}]}$. If $q \leq p$ and $\bar{a}$ indirectly approximates q , then $q \leq p^{[\bar{a}]}$.
- (2) If $q \leq_g p$, then $q^{[\bar{a}]} \leq p^{[\bar{a}]}$.
- (3) If $q \leq p^{[\bar{a}]}$ then there is a $r \leq_g p$ such that $r^{[\bar{a}]} =^* q$.
- (4) The set $\{p^{[\bar{a}]} : \bar{a} \in \text{Pos}(g)\}$ is predense below p .
- (5) We can densely determine η_i up to n , i.e. for all $i \in I$, $n \in \omega$ the following set is dense: $\{p \in P : (\exists a \in \omega^n)(i, a) \text{ approximates } p\}$.
- (6) (Pure decision) If $D \subseteq P$ is open dense, and g indirectly approximates p , then there is an $r \leq_g p$ such that $r^{[\bar{a}]} \in D$ for all $\bar{a} \in \text{Pos}(g)$.

Proof. (1)–(3) follow from the last Facts.

(4) If g indirectly approximates p and $q \leq p$, then there is a h stronger than g approximating q . Let $\bar{b} \in \text{Pos}(h)$ and $\bar{a} = \bar{b} \upharpoonright g \in \text{Pos}(g)$. Then $q^{[\bar{b}]} \leq q$, $p^{[\bar{a}]}$.

(5) Let h approximates p such that $\text{minlength}_{[i]}(h) > n$. Let $\bar{a} \in \text{Pos}(h)$. Then (i, a_i) indirectly approximates $p^{[\bar{a}]} \leq p$. By 3.12(1) we can find a $q =^* p$ such that a_i approximates q .

(6) Let $\text{Pos}(\mathfrak{g}) = \{\bar{a}_0, \dots, \bar{a}_l\}$. Pick $q_0 \leq p^{[\bar{a}_0]}$ in D , and $r_0 \leq_{\mathfrak{g}} p$ as in (3). So $r_0^{[\bar{a}_0]} \in D$. Pick $q_1 \leq r_0^{[\bar{a}_1]}$ in D and $r_1 \leq_{\mathfrak{g}} r_0$ as above, etc. Then r_l has the required property. $\square$

We now list some trivial properties of P regarding restriction:

Definition 3.21. For $i \in I^\infty$ we define $P_i := P_{<i} := \{p \in P : \text{dom}(p) \subseteq I_{<i}\}$, and analogously for $P_{\leq i}$.

Facts 3.22. (Restriction) Assume $p, q \in P$ and $i, j \in I^\infty$.

- If $\text{dom}(q) \supseteq \text{dom}(p)$, $q \restriction \text{dom}(p) = p$ and $\mathfrak{g}$ approximates $\mathfrak{g}$, then $q \leq_{\mathfrak{g}} p$.
- $P = P_\infty$, $p \restriction I_{<i} \in P_{<i}$ and $p \leq p \restriction I_{<i}$.
- If $p' \leq p$ then $p' \restriction I_{<i} \leq p \restriction I_{<i}$. If $p \in P_{<i}$ then $p \restriction I_{<i} = p$.
- Let $q \in I_{<i}$, $q \leq p \restriction I_{<i}$. Define $q \wedge p := q \cup p \restriction I_{\geq i}$. Then $q \wedge p \in P$ is the weakest condition stronger than both q and p .
- $p \restriction I_{<i}$ is a reduction of p (i.e. $r' \in P_{<i}$ and $r' \leq p \restriction I_{<i}$ implies $r' \parallel p$).
- If $i \leq j$ then $P_{<i} < P_{<j}$ (i.e. $P_{<i}$ is a complete subforcing of $P_{<j}$).
- If $p \restriction I_{<i} \parallel q \restriction I_{<i}$ and $\text{dom}(p) \cap \text{dom}(q) \subseteq I_{<i}$, then $p \parallel q$.
- Similar facts hold for $P_{\leq i}$. E.g. if $i < j$, then $P_{\leq i} < P_{<j}$.

Definition 3.23. Assume that $i \leq j \in I^\infty$ and that G_j is a $P_{<j}$ -generic filter over V .

- The filter induced by the complete embedding $P_{<i} < P_{<j}$ is called G_i , and $V[G_i]_{P_{<i}}$ is called V_i . The canonical Q_i -generic filter over V_i is called $G(i)$. Analogously we define $V_{\leq i}$ and $G_{\leq i}$ (which will turn out to be $V_i[G(i)]$ and $G_i * G(i)$).
- In V_j or $V_{\leq i}$ we define η_i to be the union of all $t \in \omega^{<\omega}$ such that (i, t) is an approximation of p for some $p \in G_j$ (or $G_{\leq i}$).

Lemma 3.24. Let i, j, G_j be as above, $p \in G_j$, and set $\bar{\eta} = (\eta_i)_{i \leq j}$.

- (1) η_i is a well-defined real.
- (2) If $\mathfrak{g}$ indirectly approximates p , then $\bar{\eta}$ is compatible with $\mathfrak{g}$.
- (3) $\{\eta_i\} = \bigcap \{\lim B_i^p(\bar{\eta}) : p \in G_j, i \in \text{dom}(p)\}$
- (4) If $p \Vdash (\forall i \in \text{dom}(q)) \eta_i \in \lim(B_i^q(\eta))$, then $q \leq p$.
- (5) $q \in G$ iff $\eta_i \in \lim(B_i^q(\bar{\eta}))$ for all $i \in \text{dom}(q)$.

Proof. (1) By 3.20(5), the set of conditions q such that for some t of length n (i, t) is an approximation of q is dense. Therefore η_i is infinite. Also, If $s \perp t$, (i, t) is an approximation of q , and (i, s) is an approximation of q' , then q and q' are incompatible. This shows that η_i is indeed a real.

(2) According to 3.20(4), the set $\{p^{[\bar{a}]} : \bar{a} \in \text{Pos}(\mathfrak{g})\}$ is predense below p . Let $\bar{a}$ be such that $p^{[\bar{a}]} \in G$. Any $q \in G$ that is stronger than $p^{[\bar{a}]}$ and decides η_i up to the length of a_i forces that $\eta_i \supset a_i$. So $\bar{\eta}$ is compatible with $\bar{a}$ and therefore with $\mathfrak{g}$.

(3) Let $n \in \omega$. We have to show that $\eta_i \restriction n \in B_i^p(\bar{\eta})$. First pick an approximation $\mathfrak{g}$ of p with $\text{minlength}_{\eta_i}(\mathfrak{g}) \geq n$. We already know that $\bar{\eta}$ is compatible with $\mathfrak{g}$, in particular η_i is compatible with $\mathfrak{g}(i)(\bar{\eta})$. And $\mathfrak{g}(i)(\bar{\eta})$ is a front in $B_i^p(\bar{\eta})$, since $\mathfrak{g}$ approximates p .

(4) Let $\mathfrak{g}$ approximate p and $\mathfrak{h}$ approximate q such that $\text{dom}(\mathfrak{h}) \supseteq \text{dom}(\mathfrak{g})$ and the length of $\mathfrak{h}$ is sufficiently large on $\text{dom}(\mathfrak{g})$. Then $\mathfrak{h}$ must be stronger than $\mathfrak{g}$.

(5) follows from (4). $\square$

As immediate consequence of fusion and pure decision we get:

Theorem 3.25. (1) P is ω^ω -bounding. For every p and P -name τ for an ω -sequence of ordinals there is a $p' \leq p$ such that p' reads τ continuously.

(2) Assume the cofinality⁸ of I is $\geq \aleph_1$, G is P -generic over V and $r \in \mathbb{R}^{V[G]}$. Then there is an $i \in I$ such that $r \in \mathbb{R}^{V_i}$.

⁸We always mean the “upwards cofinality”, i.e. the minimal size of an upwards cofinal subset. $A \subset I$ is upwards cofinal if for every $i \in I$ there is an $a \in A$ such that $a \geq i$.

- (3) P is proper.
- (4) P forces that η_i is a Q_i -generic real over V_i .
- (5) If $I = I_1 + I_2$, then $\text{nwf-lim}_I(Q_i) \cong \text{nwf-lim}_{I_1}(Q_i) * \text{nwf-lim}_{I_2}(Q_i)$.
- (6) If $I = \sum_{\beta \in \epsilon} J_\beta$ is the concatenation of the orders J_β along the ordinal ϵ , then $\text{nwf-lim}_I(Q_i)$ is equivalent to the countable support limit $(P_\beta, \mathcal{Q}'_\beta)_{\beta \in \epsilon}$ of the forcings $\mathcal{Q}'_\beta := \text{nwf-lim}_{J_\beta}(Q_i)$.
- (7) If I is well-founded, then $\text{nwf-lim}_I(Q_i)$ is the countable support limit of the Q_i .

Proof. (1) Fix for every countable subset J of I an enumeration $\{j_m : m \in \omega\}$, and denote $\{j_m : m \in n\}$ by $\text{first}(n, J)$.

Assume τ is a name of a real and $p \in P$. We have to show that there is a $p_\omega \leq p$ and an $f \in \omega^\omega$ such that $p_\omega \Vdash \tau(n) < f(n)$. Let $p_0 \leq p$, $f(0) \in \omega$ be such that $p_0 \Vdash \tau(0) = f(0)$, and let g_0 approximate p_0 . Assume that g_n and p_n are already defined. We define $p_{n+1} \leq_{g_n} p_n$, $f(n)$ and g_{n+1} the following way: Let $p_{n+1} \leq_{g_n} p_n$ be such that $p_{n+1}^{[\bar{a}]}$ decides $\tau(n)$ for every $\bar{a} \in \text{Pos}(g_n)$, see 3.20(6). Let $f(n)$ be the maximum of the possible values for $\tau(n)$. Let g_{n+1} be a p_{n+1} -approximation stronger than g_n which is n -dense at every $i \in \text{first}(n, \text{dom}(p_1)) \cup \dots \cup \text{first}(n, \text{dom}(p_n))$. Then the sequence $(p_n)_{n \in \omega}$ satisfies the conditions for fusion 3.11 so there is a $p_\omega \leq p_n$. Clearly, $p_\omega \Vdash \tau(n) \leq f(n)$.

(2) The p_ω above completely determines τ , so if $p_\omega \in P_{<i}$, then $p_\omega \Vdash_P \tau \in V_i$.

(3) is very similar to the above: Assume that $N < H(\chi)$ and $p_0 \in N$. Let $\{D_m : m \in \omega\}$ enumerate the dense sets in N . Assume $p_n, g_n \in N$ are already defined. Find (in N) $p_{n+1} \leq_{g_n} p_n$ such that $p_{n+1}^{[\bar{a}]} \in D_n$ for all $\bar{a} \in \text{Pos}(g_n)$, and pick $g_{n+1} \in N$ big enough. Then we can (in V) fuse this sequence into a $p_\omega \in P$. Note that $\text{dom}(p_\omega) \subseteq N \cap I$. If G is P -generic over V and $p_\omega \in G$, then $p_n \in G$ and $\{p_n^{[\bar{a}]} : \bar{a} \in \text{Pos}(g_n)\}$ is predense below p_n , so some $p_n^{[\bar{a}]} \in G$, and $p_n^{[\bar{a}]} \in D_n \cap N$.

(4) is a special case of (5): Set $I_1 := I_{<i}$ and $I_2 := \{i\}$. So η_i is V_i -generic in $V_{\leq i}$ and therefore in V_∞ as well.

(5) Set $P := \text{nwf-lim}_I(Q_i)$, $P_1 := \text{nwf-lim}_{I_1}(Q_i)$, $P_2 := \text{nwf-lim}_{I_2}(Q_i)$.

There is a natural map $\phi : p \mapsto (p_1, p_2)$ from P to $P_1 * P_2$: $p_1 := p \restriction I_1$, and p_2 is defined by $\text{dom}(p_2) := \text{dom}(p) \setminus I_1$ and $B_i^{p_2}(\bar{\eta}) := B_i^p((\eta_i)_{i \in I_1} \bar{\eta})$.

It is clear that ϕ preserves $\leq$. We claim that it is dense and preserves $\perp$. Assume $\phi(p) = (p_1, p_2)$, $\phi(q) = (q_1, q_2)$, and $(r_1, r_2) \leq (p_1, p_2), (q_1, q_2)$. We have to find a $r' \leq_P p, q$ such that $\phi(r') \leq (r_1, r_2)$.

r_1 forces that p_2, q_2 and r_2 correspond to approximating sequences $(g_n^p), (g_n^q)$ and $(g_n^{r'})$. As in (1) we can find an $r'_1 \leq r_1$ with an approximating sequence (h_n) such that h_n decides g_n^i (for $i \in \{p, q, r\}$) in a way such that $g_n^{r'}$ is stronger than both g_n^p and g_n^q . Then we can concatenate (h_n) with the $(g_n^{r'})$ to an approximating sequence to some $r' \in P$. Then $r' \leq p, q$ and $\phi(r') \leq (r_1, r_2)$.

(6) By induction on ϵ . The successor step follows from (5). Let $\text{cf}(\epsilon) > \omega$. Then the nwf-limit as well as the cs-limit are just the unions of the smaller limits, and therefore equal by induction. If $\text{cf}(\epsilon) = \omega$, then the nwf-limit as well as the cs-limit are the full inverse limits of the iteration system, and therefore again equal by induction.

(7) follows from (6). $\square$

We will also use the following fact:

Lemma 3.26. *If $q \restriction I_i$ reads $\mathcal{S} \in Q$ continuously and $q \Vdash \eta_i \in \mathcal{S}$, then $q \restriction I_i$ forces that $B_i^q(\bar{\eta}) \leq_Q \mathcal{S}$.*

Proof. Assume otherwise. Then there is an approximation g of $p := q \restriction I_i$, an $\bar{a} \in \text{Pos}(g)$ and a $t \in T_{\max}^i$ such that $p^{[\bar{a}]}$ forces t to be in $B_i^q(\bar{\eta})$ but not in $\mathcal{S}$. Let $\bar{a}^+$ be $\bar{a} \frown t$. Then $\bar{a}^+$ is a possible value of some approximation of q , and $q^{[\bar{a}^+]}$ forces that $\eta_i \notin \mathcal{S}$, a contradiction. $\square$

Remark. The iteration technique defined here also works for larger classes of forcings, e.g. for the tree forcings $\mathbb{Q}_0^{\text{tree}}$ of [6] mentioned already. If we assume additional properties

such as bigness and halving, we could also use $\lim\text{-inf}$ forcings. It is also possible to extend the construction to non-total orders, or to allow $T_{\max}^i, \mu^i$ to be $P_{<i}$ -names.

4. $\mathbb{T}$ $\mathbb{I}^c$

In this section we define (to a $\lim\text{-sup}$ forcing Q as in section 2) ideals $\mathbb{I}$ and $\mathbb{I}^c$ (the $< 2^{\aleph_0}$ -closure of $\mathbb{I}$). These ideals will generally not be ccc. We define what we mean by “ X is measurable with respect to $\mathbb{I}^c$ ”. The main theorem of this paper is that for certain Q we can force measurability for all definable X without using an inaccessible or amalgamation of forcings.

Definition 4.1. • The ideal $\mathbb{I}$ on the reals is defined by: $X \in \mathbb{I}$ if for all $S \in Q$ there is a $T \leq S$ such that $X \cap \lim(T) = \emptyset$.
 • $\mathbb{I}^c$ is the $< 2^{\aleph_0}$ -closure of $\mathbb{I}$.
 • X is null if $X \in \mathbb{I}^c$; X is positive if $X \notin \mathbb{I}^c$; X has measure 1 if $2^\omega \setminus X \in \mathbb{I}^c$.

Notes. • Of course the notion null etc. depend on the forcing Q , so it might be more exact to use notation such as Q -null or $(T_{\max}, \mu)$ -null. In this paper this is not necessary, since we will always use a fixed Q .
 • We use the words “null”, “positive” and “measure 1” although the ideals $\mathbb{I}$ and $\mathbb{I}^c$ are not related to a measure (they are not even ccc).
 • If CH holds, then $\mathbb{I}^c = \mathbb{I}$.
 • $\mathbb{I}$ is always nontrivial (i.e. $V \notin \mathbb{I}$), but this is not clear for $\mathbb{I}^c$.

$F : Q \rightarrow Q$ is a witness for $X \in \mathbb{I}$ if $F(S) \leq S$ and $X \cap \lim(F(S)) = \emptyset$ for all $S \in Q$.
 So every $X \in \mathbb{I}$ is contained in a set $\bigcap \{\omega^\omega \setminus \lim(F(S)) : S \in Q\}$.⁹

Lemma 4.2. $\mathbb{I}$ is a non-trivial σ -ideal.

Proof. This follows from fusion: Assume $X_i \in \mathbb{I}$ ($i \in \omega$) and $S = S_0 \in Q$. Pick any front $F_0 \in S_0$, so $S_0 = \bigcup_{t \in F_0} S_{0,t}^{[t]}$. For each $t \in F_0$ pick an $S_{1,t} \leq S_{0,t}^{[t]}$ such that $\lim(S_{1,t}) \cap X_1 = \emptyset$. Set $S_1 := \bigcup_{t \in F_0} S_{1,t}$. So $S_1 \in Q$, and F_0 is a front in S_1 . Pick a 1-dense front F_1 in S_1 (purely) stronger than F_0 . Iterate the construction. Fusion produces a $T < S$ such that $\lim(T) \cap X_i = \emptyset$ for all $i \in \mathbb{N}$. $\square$

For example, if Q is Sacks forcing, then $\mathbb{I}$ is called Marczewski ideal. $X \in \mathbb{I}$ iff in every perfect set A there is a perfect subset A' of A such that $A' \cap X = \emptyset$. So if X is Borel (or if X has the perfect set property, e.g. X is Σ_1^1), then $X \in \mathbb{I}$ iff X is countable. $\mathbb{I}$ is not a ccc ideal: For $A \subseteq \omega$, set

$$X_A := \{f \in 2^\omega : (\forall n \notin A) f(n) = 0\}.$$

Clearly $X_A \cap X_B = X_{A \cap B}$, and $|X_A| = 2^{|A|}$. So if $\{A_i : i \in 2^{\aleph_0}\}$ is an almost disjoint family, then $\{X_{A_i}\}$ is a family of closed sets not in $\mathbb{I}$ such that $X_{A_i} \cap X_{A_j}$ is finite for $i \neq j$.

For a Borel ccc ideal I , “ $X \subseteq \mathbb{R}$ is measurable” can be defined by “there is a Borel set A such that $A \Delta X \in I$ ”. (Usually the basis of the ideal is simpler, e.g. one can use open sets instead of Borel sets for meager, or G_δ sets for Lebesgue-null.) Equivalently, X is measurable iff for every I -positive Borel set A there is an I -positive Borel set $B \subseteq A$ such that either $B \cap X \in I$ or $B \setminus X \in I$. For non-ccc ideals that do not live on the Borel sets, this second notion is usually the right one to generalize.

In our case we use the following definition of measurability:

Definition 4.3. • $X \subseteq \mathbb{R}$ is strongly measurable if for every $T \in Q$ there is an $S \leq_Q T$ such that either $\lim(S) \cap X \in \mathbb{I}$ or $\lim(S) \setminus X \in \mathbb{I}$.
 • $X \subseteq \mathbb{R}$ is measurable if for every $T \in Q$ there is an $S \leq_Q T$ such that either $\lim(S) \cap X \in \mathbb{I}^c$ or $\lim(S) \setminus X \in \mathbb{I}^c$.

⁹Note that this is not a countable intersection.

Again, it would be more exact (but it is not necessary here) to say “ $(T_{\max}, \mu)$ -measurable” instead of “measurable”.

Clearly strong measurability implies measurability.

Lemma 4.4. *The family of (strongly) measurable sets contains the Borel sets and is closed under complements and countable unions.*

Proof. Closure under complement is trivial.

Every closed set is strongly measurable: Let $X = \lim(T')$ be closed and $T \in Q$. If there is a $t \in T \setminus T'$ then $S := T^{[t]}$ satisfies $\lim(S) \cap X = \emptyset$. Otherwise $T \subseteq T'$ and $S := T$ satisfies $\lim(S) \setminus X = \emptyset$.

Assume that $(X_i)_{i \in \omega}$ is a sequence of measurable sets and that $T \in Q$. If for some $i \in \omega$ there is an $S \leq T$ such that $\lim(S_i) \setminus X_i \in \mathbb{I}^c$ then the same obviously holds for $\bigcup_{i \in \omega} X_i$. So assume that for all $i \in \omega$ and $T' \leq T$ there is an $S \leq T'$ such that $\lim(S) \cap X_i \in \mathbb{I}^c$. Now repeat the proof of 4.2.

The same proof shows that the *strongly* measurable sets are closed under countable unions as well. $\square$

$\mathbb{I}^c$ could be trivial. However, if it is “everywhere nontrivial”, then $\mathbb{I}^c$ and $\mathbb{I}$ are the same on Borel sets:

Lemma 4.5. *Assume that $\lim(S) \notin \mathbb{I}^c$ for all $S \in Q$. Then $\mathbb{I}^c$ and $\mathbb{I}$ agree on strongly measurable sets. I.e. if X is strongly measurable and $X \in \mathbb{I}^c$, then $X \in \mathbb{I}$.*

In particular, in this case every Borel set is in $\mathbb{I}^c$ iff it is in $\mathbb{I}$.

Proof. For every $T \in Q$ there is an $S \leq_Q T$ such that $\lim(S) \cap X \in \mathbb{I}$: Otherwise $\lim(S) \setminus X \in \mathbb{I} \subseteq \mathbb{I}^c$, a contradiction to $X \in \mathbb{I}^c$ and $\lim(S) \notin \mathbb{I}^c$. So by the definition of $\mathbb{I}$ there is a $S' \leq_Q S \leq_Q T$ such that $\lim(S') \cap X = \emptyset$. So $X \in \mathbb{I}$. $\square$

Fact 4.6. *For Borel codes B , the statement “ $B \in \mathbb{I}$ ” is $\mathbb{I}_2^1$ and therefore invariant under forcing.*

On the other hand, since $\mathbb{I}$ is not a Borel ideal (i.e. not every $X \in \mathbb{I}$ is contained in a Borel set $B \in \mathbb{I}$), there is no reason why $X \in \mathbb{I}$ should be upwards absolute between universes.

The main theorem of this paper states that (using an artificially sweet forcing P) we can force all definable sets to be measurable:

Theorem 4.7. *Assume CH and that Q satisfies the Ramsey property 6.4.*

Then there is a proper, $\aleph_2$ -cc, ω^ω -bounding p.o. P forcing that every set of reals which is (first-order) definable using a parameter in $L(\mathbb{R})$ is measurable.

P will be the non-wellfounded iteration of Q along an order I defined in the next section.

We already know the following (use 2.8):

Lemma 4.8. *If I has cofinality $\geq \aleph_1$ and $i \in I$ then $\Vdash_P V_i \in \mathbb{I}$.*

Proof. If $T \in V[G_P]$ then $T \in V_j$ for some $i < j < \infty$ because of 3.25(2). So in $V_{\leq j}$ there is an $S < T$ such that $\lim(S) \cap V_i = \emptyset$ (in $V_{\leq j}$ and $V[G_P]$ as well, according to 2.8). $\square$

For later reference, we will reformulate the definition of $\mathbb{I}$: If $S \in Q$, $X \subseteq Q$, $T \in X$ and $T' \leq_Q S, T$, then $\lim(T') \cap (2^\omega \setminus \bigcup_{T \in X} \lim(T)) \subseteq \lim(T') \setminus \lim(T) = \emptyset$. So we get:

Lemma 4.9. *If $X \subseteq Q$ is predense then $\bigcup_{T \in X} \lim(T)$ is of measure 1 with respect to $\mathbb{I}$.*

5. A

In this section we assume CH. We will construct an order I and define P to be the nwf-limit of Q along I . I is ω_2 -like,¹⁰ has a cofinal sequence j_α ($\alpha \in \omega_2$) and many automorphisms. We show that these properties imply that P forces the following:

¹⁰ I is ω_2 -like if $|I_{<i}| < \aleph_2$ for all $i \in I$ and $|I| = \aleph_2$.

- $2^{\aleph_0} = \aleph_2$,
- $\mathbb{I}^c$ is nontrivial (and moreover $\lim(S) \notin \mathbb{I}^c$ for all $S \in Q$),
- for every definable set X , “locally” either all or no η_{j_δ} are in X and
- $\{\eta_{j_\delta} : \delta \in \omega_2\}$ is of measure 1 in $\{\eta_i : i \in I\}$.

In the next section it will be shown that the set $\{\eta_i : i \in I\}$ is of measure 1, which will finish the proof of the main theorem.

Lemma 5.1. *Assume CH holds and I is ω_2 -like. Then*

- (1) P has the $\aleph_2$ -cc (and therefore preserves all cofinalities).
- (2) $P_{<i} \Vdash CH$ for each $i \in I$ and $P \Vdash 2^{\aleph_0} = \aleph_2$.

Proof. (1) If $|I_{<i}| \leq 2^{\aleph_0}$ then $|P_{<i}| \leq 2^{\aleph_0}$: There are at most $|I_{<i}|^{\aleph_0} \leq 2^{\aleph_0}$ many countable subsets of $|I_{<i}|$. For each $p \in P_{<i}$ with a fixed domain and each $j \in \text{dom}(p)$ there are $2^{\aleph_0}$ many possibilities for Dom_j^p and $2^{\aleph_0}$ many possibilities for the Borel definition B_j^p .

If CH holds, then the usual delta system lemma applies: If $A \subseteq P$ is a maximal antichain of size $\aleph_2$ then without loss of generality the domains of $p \in A$ form a delta system (i.e. there is a countable $x \subseteq I$ such that $\text{dom}(p_1) \cap \text{dom}(p_2) = x$ for all $p_1 \neq p_2 \in A$). Since I is ω_2 -like, x cannot be cofinal. Let i be an upper bound of x . Without loss of generality $p_1 \restriction I_{<i} = p_2 \restriction I_{<i}$ for $p_1 \neq p_2 \in A$ (since there are only $\aleph_1$ many elements of $P_{<i}$). But then $p_1 \parallel p_2$ by 3.22.

Proper and $\aleph_2$ -cc imply preservation of all cofinalities and cardinalities.

(2) Let G be P -generic over V . Then the reals in $V[G_P]$ are the union of the reals in $V[G_{P_{<i}}]$, and every real in $V[G_{P_{<i}}]$ is decided by (i.e. read continuously from) a condition $p \in P_{<i}$. Since $|P_{<i}| = (2^{\aleph_0})^V = \aleph_1$, there are at most $\aleph_2$ many reals in $V[G_{P_{<i}}]$. And $\eta_i \notin V[G_{P_i}]$, so in particular $\eta_{i_1} \neq \eta_{i_2}$ for $i_1 \neq i_2$. $\square$

The following is well known:

Lemma 5.2. *If CH holds, then there is an $\aleph_1$ saturated¹¹ linear order $\tilde{I}$ of size $\aleph_1$, and all such orders are isomorphic.*

Proof. Induction of length ω_1 : Assume at stage α we have a linear order L_α of size $\omega_1 = 2^{\aleph_0}$. List all the $(\omega_1$ many) gaps of the types (a, b) , where $a, b \in \{1, \aleph_0, \pm\infty\}$, and add points to fill these gaps. At limits, take the union. Then at stage ω_1 we get a saturated order.

Uniqueness is proven by the standard back and forth argument. $\square$

Definition 5.3. Let $\mathfrak{S}$ be the set of $0 < \alpha < \omega_2$ such that $\text{cf}(\alpha) \in \{1, \omega_1\}$. Note that $\mathfrak{S} \subseteq \omega_2$ is stationary.

We will now define the order I along which we iterate. (We do this assuming CH.)

Given $\tilde{I}$ as above, let I be the following order:

$$\underbrace{\tilde{I}}_0 + \underbrace{\{j_1\} + \tilde{I} + \cdots}_1 + \underbrace{\tilde{I}}_\omega + \underbrace{\{j_{\omega+1}\} + \tilde{I} + \cdots}_{\omega+1} + \underbrace{\{j_{\omega_1}\} + \tilde{I} + \cdots}_{\omega_1}$$

So at stages $\alpha \in \mathfrak{S}$, we add $\{i\} + \tilde{I}$, in other stages we add just $\tilde{I}$.

- Facts 5.4.**
- I is ω_2 -like,
 - $(j_\alpha)_{\alpha \in \mathfrak{S}}$ is an increasing (and therefore cofinal) continuous sequence in I , and
 - every j_α has cofinality $\aleph_1$ in I .

Continuous means that $j_\delta = \sup(j_\alpha : \alpha \in \mathfrak{S}, \alpha < \delta)$ whenever $\delta = \sup(\mathfrak{S} \cap \delta) \in \mathfrak{S}$ (which is equivalent to $\text{cf}(\delta) = \omega_1$).

Note. We could just as well define j_α for α with cofinality ω_1 only, or for all $\alpha \in \omega_2$ (and require continuity for points of cofinality ω_1 only). All these versions are equivalent by simple relabeling, cf. the beginning of the proof of 5.7.

¹¹A linear order $\tilde{I}$ is $\aleph_1$ saturated if there are no gaps of type (a, b) , where $a, b \in \{1, \aleph_0, \pm\infty\}$.

Definition 5.5. We set $Q_i = Q$ for all $i \in I$ and let P be the nwf-iteration of Q_i along I .

We will use the notation $I_\alpha, P_\alpha, V_\alpha$ and η_α for $I_{<j_\alpha}, P_{<j_\alpha}, V_{j_\alpha}$ and η_{j_α} .

We set $V_{\omega_2} := V_\infty = V[G_P]$.

Lemma 5.6. (CH) Let $S_0 \subseteq \mathfrak{S}$ be stationary. P forces the following:

- (1) $\{\eta_\delta : \delta \in S\} \notin \mathbb{I}^c$ for every stationary $S \subseteq \mathfrak{S}$, and
- (2) $\{\eta_\delta : \delta \in S_0\} \cap \lim(T_0) \notin \mathbb{I}^c$ for every $T_0 \in Q$.

This lemma implies that in V_{ω_2} the assumption of Lemma 4.5 is satisfied (i.e. that $\mathbb{I}^c$ is “everywhere nontrivial”). Note that this lemma holds for all I satisfying 5.4.

Proof. (1) Assume otherwise, i.e. there are P -names $\tilde{F}_\zeta$ ($\zeta \in \omega_1$) for functions from Q to Q and $\mathfrak{S}$ for a stationary set such that $p_0 \in P$ forces

$$\tilde{F}_\zeta(T) \leq T \text{ and } (\forall \delta \in S) (\exists \zeta \in \omega_1) (\forall T \in Q) \eta_\delta \notin \lim(\tilde{F}_\zeta(T)).$$

P forces that for each $\alpha \in \mathfrak{S}$ there is a $\beta \in \mathfrak{S}$ such that $\tilde{F}_\zeta(T) \in Q^{V_\beta}$ for all $T \in Q^{V_\alpha}$ and $\zeta \in \omega_1$. We need something slightly stronger: For every name $\tilde{T}$ for an element of Q^{V_α} and $\zeta \in \omega_1$ there is a maximal antichain $A \subset P$ such that for every $q \in A$ there is a P -name $\tilde{T}'_q$ such that q forces $\tilde{F}_\zeta(\tilde{T}) = \tilde{T}'_q$ and q continuously reads $\tilde{T}'_q$. So if $q \in G_P$ and β is bigger than $\text{dom}(q)$,¹² then V_β not only contains $T'_q = \tilde{F}_\zeta(T)$, but also knows that T'_q will be $\tilde{F}_\zeta(T)$ in V_{ω_2} .

Define $f^-(\alpha)$ to be the smallest β which is bigger than $\text{dom}(q)$ for every $a \in A$, where A is an antichain for some $\tilde{T}$ and $\zeta \in \omega_1$ as above. P is $\aleph_2$ -cc, every $q \in A$ has countable domain, and there are only $\aleph_1$ many reals in V_α . So $f^-(\alpha) < \omega_1$, and we can define $f(\alpha)$ to be the smallest $\beta \in \mathfrak{S}$ that is larger or equal to $\max(\alpha, f^-(\alpha))$.

If $\text{cf}(\alpha) = \omega_1$, then $f(\alpha)$ is the supremum of $\{f(\gamma) : \gamma \in \mathfrak{S} \cap \alpha\}$, since the reals in V_α are the union of the reals in V_γ . So f is continuous.

Then P forces the following: Since $\mathfrak{S}$ is stationary, there is a $\beta \in \mathfrak{S}$ such that $f(\beta) = (\beta)$. V_β can calculate every $\tilde{F}_\zeta$, and $\tilde{F}_\zeta''Q$ is dense in Q . Since η_β is a Q -generic real over V_β , there is (for every $\zeta \in \omega_1$) a $T \in Q^{V_\beta}$ such that $\eta_\beta \in \lim(\tilde{F}_\zeta(T))$, a contradiction.

(2): We can assume that $T_0 \in V$. Again, choose names $\tilde{F}_\zeta$ as above, and assume that $p_0 \in P$ forces that

$$\tilde{F}_\zeta(T) \leq T \text{ and } (\forall \delta \in S_0) (\exists \zeta \in \omega_1) (\forall T \in Q) \eta_\delta \notin \lim(\tilde{F}_\zeta(T)) \cap \lim(T_0).$$

Define f as above, so there is a $\beta > \text{dom}(p)$ such that $\beta \in S_0$ and $f(\beta) = \beta$. So the same argument proves that p_0 forces that $\eta_\beta \notin \lim(T_0)$, a contradiction. $\square$

We also get the following:

Lemma 5.7. (CH) For every $C \subseteq \omega_2$ club, P forces the following:

$$\{\eta_i : i \in I\} \setminus \{\eta_\alpha : \alpha \in \mathfrak{S} \cap C\} \in \mathbb{I}^c.$$

Again, this lemma applies to all I satisfying 5.4.

Proof. We can assume that $C = \omega_2$, since we can just relabel the sequence $\{j_\alpha : \alpha \in \mathfrak{S} \cap C\}$: Set $j'_\alpha := j_\beta$, where β is the α -th element of $C \cap \mathfrak{S}$. Then $(j'_\alpha)_{\alpha \in \mathfrak{S}}$ satisfies 5.4 as well.


Recall Definition 2.9 of $Q_{A_\gamma}^f$ and D_f^{spl} (for $f : \omega \rightarrow \omega$ increasing and $r \in 2^\omega$). Enumerate all increasing $f : \omega \rightarrow \omega$ in V as f_ζ ($\zeta \in \omega_1$). (CH holds in V .)

Claim: In V , we can find P_α -names $\tilde{T}_\alpha^\zeta$ ($\zeta \in \omega_1, \alpha < \omega_2$ successor) for elements of Q such that the following is forced by P (cf. Figure 5):

- (1) The set $\{\tilde{T}_{\alpha+1}^\zeta : \alpha < \omega_2\} \subseteq Q$ is dense for all $\zeta \in \omega_1$.
- (2) $\tilde{T}_\alpha^\zeta \in D_{f_\zeta}^{\text{spl}}$ (in V_α or equivalently in V_{ω_2}).¹³
- (3) If $\beta < \alpha$ is a successor, then $\tilde{T}_\alpha^\zeta$ has no branch in V_β , and for all $i < j_\alpha$ there is a ζ_0 such that $\tilde{T}_\alpha^\zeta$ has no branch in V_i for all $\zeta \geq \zeta_0$.

¹²More formally: if $j_\beta > i$ for all $i \in \text{dom}(q)$.

¹³Recall 2.9 and 2.10.



F 5.

Proof of the claim: Pick for all $\alpha+1$ a function $\phi_{\alpha+1} : \omega_1 \rightarrow I_{j_{\alpha+1}} \setminus I_{j_\alpha}$ which is increasing and cofinal. Also pick an enumeration $(\zeta_{\alpha+1})_{\alpha \in \omega_2}$ such that ζ_α is a P_α -name and P forces that $Q = \{\zeta_{\alpha+1} : \alpha \in \omega_2\}$. (This is possible since P forces that $Q^{V_{\omega_2}} = \bigcup Q^{V_\alpha}$, cf. 3.25(2).)

To find T_α^ζ (α successor) note that P_α forces that we can perform the following construction in V_α :

First pick an $S' \leq \zeta_\alpha$ such that $S' \in D_{f_\zeta}^{\text{spl}}$ (cf. 2.10(7)). $\text{cf}(j_\alpha) = \aleph_1$, so $S' \in V_i$ for some $i < j_\alpha$. Pick some i' bigger than $\max(i, \phi_\alpha(\zeta))$ and smaller than j_α . There is a real $r \in V_\alpha \setminus V_{i'}$ (e.g. $\eta_{i'}$). Therefore there is a $T_\alpha^\zeta < S'$ such that $\lim(T_\alpha^\zeta) \cap V_{i'} = \emptyset$ (in V_α and V_{ω_2} as well, cf. 2.8). Let T_α^ζ be a P_α -name for T_α^ζ .

The T_α^ζ constructed this way satisfy the claim: (1): $T_\alpha^\zeta \leq \zeta_\alpha$, (2): $D_{f_\zeta}^{\text{spl}}$ is open dense and absolute, (3): pick ζ_0 such that $\phi_\alpha(\zeta_0) > i$. This ends the proof of the claim.

From now on assume G is P -generic over V . We work in V_{ω_2} and set $T_\alpha^\zeta := T_\alpha^\zeta[G]$. So if $i \in I$ then the sequence $(T_{\alpha+1}^\zeta)_{j_{\alpha+1} < i, \zeta \in \omega_1}$ is in V_i .

For all $\zeta \in \omega_1$, $X_\zeta := \bigcup_{\alpha+1 < \omega_2} \lim(T_{\alpha+1}^\zeta)$ is of measure 1 with respect to $\mathbb{I}$ (cf. 4.9). So the set $Y := \bigcap_{\zeta \in \omega_1} X_\zeta$ is of measure 1. It is enough to show that

$$\{\eta_i : i \in I\} \setminus \{\eta_\alpha : \alpha \in \mathfrak{S}\} \cap Y = \emptyset.$$

Assume towards a contradiction that some η_i is in Y and $\eta_i \neq \eta_\alpha$ for all $\alpha \in \mathfrak{S}$.

Let $\alpha \in \mathfrak{S}$ be minimal such that $\eta_i \in V_\alpha$ (i.e. $i < j_\alpha$, cf. Figure 5). So α is a successor (but not necessarily a successor of a $\beta \in \mathfrak{S}$), and $i > j_\beta$ for all $\beta \in \mathfrak{S} \cap \alpha$. So according to (3) there is a ζ_0 such that $\eta_i \notin \lim(T_{\gamma+1}^\zeta)$ for all $\zeta > \zeta_0$ and all $\gamma+1 \geq \alpha$.

So we know the following: $\eta_i \in Y$, i.e.

$$\eta_i \in \bigcup_{\gamma+1 < \omega_2} \lim(T_{\gamma+1}^\zeta) \quad \text{for all } \zeta \in \omega_1.$$


But

$$\eta_i \notin \bigcup_{\alpha \leq \gamma+1 < \omega_2} \lim(T_{\gamma+1}^\zeta) \quad \text{for all } \zeta \geq \zeta_0.$$

Therefore

$$\eta_i \in \bigcup_{\gamma+1 < \alpha} \lim(T_{\gamma+1}^\zeta) \quad \text{for all } \zeta \geq \zeta_0.$$

Recall that V_i sees the sequence $(T_{\gamma+1}^\zeta)_{\gamma+1 < \alpha, \zeta \in \omega_1}$. So in V_i , some $T \in Q$ forces that for all $\zeta > \zeta_0$ there is a successor $\beta(\zeta) < \alpha$ such that $\eta_i \in \lim(T_{\beta(\zeta)}^\zeta)$. In V_i , T has full splitting for some $f_\zeta \in V$, $\zeta > \zeta_0$ (see 2.10(5), 2.11 and 3.25(1)).

F 6. An automorphism f .

Let r be a real in $V_i \setminus \bigcup_{\gamma+1 < \alpha} V_{\gamma+1}$. Pick in V_i a $T' \leq T$ such that $T' \in Q_{A_r}^{f_\zeta}$ (cf. 2.10(6)) and T' decides $\beta(\zeta)$. Then T' forces that $\eta_i \in T' \cap T_{\beta(\zeta)}^\zeta$, a contradiction to $T' \perp T_{\beta(\zeta)}^\zeta \in V_i$ (because of (2), either $T_{\beta(\zeta)}^\zeta$ is in $Q_{A_s}^{f_\zeta}$ for some old real s , or incompatible to all $Q_{A_s}^{f_\zeta}$). $\square$

We call f an automorphism if it is a $<$ -preserving bijection from I to I .

If $f : I \rightarrow I$ is an automorphism, then f defines an automorphism of P in a natural way as well (provided of course that $f(i) = j$ implies $Q_i = Q_j$, but in our case all the Q_i are the same). Also, f defines a map on all P -names, and we have: $p \Vdash \varphi(\tau)$ iff $f(p) \Vdash \varphi(f\tau)$.

If $\Vdash_P \dot{x} \in V_i$, then there is a V_i -name $\dot{\tau}$ such that $\Vdash_P \dot{x} = \dot{\tau}$. If $f \upharpoonright I_{<i}$ is the identity, then $f(\dot{\tau}) = \dot{\tau}$. So in this case $p \Vdash \phi(\dot{\tau})$ iff $f(p) \Vdash \phi(\dot{\tau})$. Also, if $f \upharpoonright \text{dom}(p) \cap I_{<i}$ is the identity then $B_i^p(\dot{\eta}) = B_{f(i)}^{f(p)}(\dot{\eta})$.

Lemma 5.8. *The following holds for I (see Figure 6): If $\alpha < \beta < \gamma < \delta \in \mathfrak{S}$, $A \subseteq I_\beta$ countable, $B \subseteq I \setminus I_\beta$ countable, then there is an automorphism f of I such that $f \upharpoonright (I_\alpha \cup A)$ is the identity, $f(j_\beta) = j_\gamma$ and $f''B > j_\delta$.*

Proof. For every $i < j \in I$, $I_{<i}$ and $\{k : i < k < j\}$ are isomorphic and also isomorphic to $\tilde{I}$ (since they are all $\aleph_1$ saturated linear orders of size $\aleph_1$). If $A \subset I$ countable, then there are $i < A < j$, and for all such i, j , $\{k : i < k < A\}$ and $\{k : A < k < j\}$ are again isomorphic to $\tilde{I}$. Also, $I_{>i}$ is isomorphic to I (since $\omega_2 \setminus \alpha$ is isomorphic to ω_2).

So assume $\alpha < \beta < \gamma \in \mathfrak{S}$, $A < i < j_\beta$ countable, $i > j_\alpha$. Then $I_{j_\beta} \setminus I_{\leq i} \cong I_{j_\gamma} \setminus I_{\leq i} \cong \tilde{I}$. Also, if $B \subset I$ is countable, $\delta \in \mathfrak{S}$ and $B > j_\beta$, then there is an $j_\beta < i < B$, and $I_i \cong I_{j_\delta} \cong \tilde{I}$, $I \setminus I_{\leq i} \cong I \setminus I_{\leq j_\delta} \cong I$. Now combine these automorphisms. $\square$

Lemma 5.9. *For $\beta \in \omega_2$ set $Y_\beta := \{\eta_\gamma : \gamma \in \mathfrak{S}, \gamma > \beta\}$. P forces the following: If X is a set of reals defined with a parameter $x \in \bigcup_{i \in I} V_i$, and if $T \in Q$, then there is an $S \leq T$ and a $\beta \in \omega_2$ such that either $\lim(S) \cap X \cap Y_\beta = \emptyset$ or $(\lim(S) \setminus X) \cap Y_\beta = \emptyset$.*

This lemma holds for all I satisfying 5.4 and 5.8.

Note that every real in V_{ω_2} is in $\bigcup_{i \in I} V_i$.

We will see in the next section that (using additional assumptions) Y_β is a measure 1 set. Then this lemma implies that X is measurable, i.e. the main theorem 4.7. Because of 5.7, it will be enough to show that the set $\{\eta_i : i \in I\}$ is of measure 1.

Proof. Assume $\dot{X} = \{r : \varphi(r, \dot{x})\}$ and fix some $\dot{T}$. Some p_0 forces that $\dot{x}$ and $\dot{T}$ are in V_α , so without loss of generality $\dot{x}, \dot{T}$ are P_α names and $\text{dom}(p_0) \subset I_\alpha$. Pick a $p_1 \leq p_0$, $p_1 \in P_\alpha$ such that p_1 continuously reads $\dot{T}$. Fix some $\beta > \alpha$. Then $p_2 := p_1 \cup \{(j_\beta, \dot{T})\}$ is an element of $P_{\leq j_\beta}$ (since $\dot{T}$ is read continuously).

Let $p \leq p_2$ decide $\varphi(\eta_\beta, \dot{x})$. Without loss of generality $p \Vdash \varphi(\eta_\beta, \dot{x})$. $p \upharpoonright I_\beta$ forces that $\dot{S} := B_{j_\beta}^p(\dot{\eta}) \leq_Q \dot{T}$ (since $p \leq p_2$).

Assume towards a contradiction that for some $q \leq p$, $\gamma \in \mathfrak{S}$ and $\gamma > \beta$

$$q \Vdash \eta_\gamma \in \lim(\dot{S}) \ \& \ \neg \varphi(\eta_\gamma, \dot{x}).$$

Note that $q \upharpoonright I_\gamma$ reads $\dot{S}$ continuously and forces that $B_{j_\gamma}^q(\dot{\eta}) \leq_Q \dot{S}$ (cf. 3.26).

Set $A := \text{dom}(p) \cap I_\beta$ and $B := \text{dom}(p) \cap I_{>j_\beta}$. Let j_δ be bigger than $\text{dom}(q)$, and let f be an automorphism of I such that $f \upharpoonright (I_\alpha \cup A)$ is the identity, $f(j_\beta) = j_\gamma$ and $f''B > \text{dom}(q)$ (cf. 5.8 or Figure 6).

$\text{dom}(f(p)) \cap \text{dom}(q) \subseteq A \cup \{j_\gamma\}$. $f(p) \upharpoonright A = p \upharpoonright A \geq q \upharpoonright A$, and $q \upharpoonright I_\gamma$ forces that

$$B_{j_\gamma}^{f(p)}(\bar{\eta}) = B_{j_\beta}^p(\bar{\eta}) = \mathcal{S} \geq_Q B_{j_\beta}^q(\bar{\eta}).$$

So $f(p)$ and q are compatible, a contradiction to $f(p) \Vdash \varphi(\eta_\gamma, \bar{x})$. $\square$

Remark. For α with cofinality ω , we could define I_α to $\bigcup_{\beta < \alpha} I_\beta$, and P_α to be the nwf-limit of the P_β ($\beta < \alpha$). Then according to Theorem 3.25, P is the c.s. limit of the iteration (P_α, R_α) , where R_α is nwf- $\lim_{\bar{I}}$ Q if $\alpha \notin \mathfrak{S}$, and nwf- $\lim_{[i] + \bar{I}}$ Q otherwise. However, we will not need this fact in this paper.

6. A -

For the proof of Theorem 4.7 it remains to be shown that $\{\eta_i : i \in I\}$ is of measure 1. For this we will need a certain Ramsey property for Q .

Definition 6.1. A subtree T of $T_{\max}$ is called (n, r) -meager if $\mu_T(t) < r$ for all $t \in T$ with length at least n .

Lemma 6.2. If T is meager for some (n, r) , then $\lim(T) \in \mathbb{I}$.

Proof. For any $S \in Q$ there is an $s \in S$ of length at least n such that $\mu_S(s) > r$. So there is an immediate successors t of s in S such that $t \notin T$. Then $\lim(S^{[t]}) \cap \lim(T) = \emptyset$. $\square$

Definition 6.3. Let M, N be natural numbers. $N \rightarrow M$ means: If

- $s_1, \dots, s_M \in T_{\max}$ such that $\text{length}(s_i) > N$,
- $t \in T_{\max}$ such that $s_i \perp t$ for $1 \leq i \leq M$,
- $A \subseteq \text{succ}(t)$ such that $\mu(A) > N$,
- $f_i : A \rightarrow T_{\max}^{[s_i]}$ for $1 \leq i \leq M$,

then there is a $B \subseteq A$ such that

- $\mu(B) > M$ and
- $\{s \in T_{\max} : (\exists i \leq M) (\exists t \in B) s \leq f_i(t)\}$ is $(N, 1/M)$ -meager.

Definition 6.4. A lim-sup tree-forcing Q is strongly non-homogeneous if μ is sub-additive¹⁴ and for all M there is an N such that $N \rightarrow M$.

There are many similar notions of bigness, see e.g. [6, 2.2].

Lemma 6.5. There is a forcing Q that is strongly non-homogeneous.

Proof. First note that it is enough to show that for each M there is an N such that $N \rightarrow^- M$, where $N \rightarrow^- M$ is defined as above but with just one s and f instead of M many. To see this, just set $K_0 := M^2$ and find K_i such that $K_{i+1} \rightarrow^- K_i$. Then $K_M \rightarrow M$. (Here we use that μ is sub-additive, since we need that the union of m many (n, r) -meager trees is $(n, r \cdot m)$ -meager.)

We will construct $T_{\max}$ and μ by induction. We define $s \triangleleft t$ by: $\text{length}(s) < \text{length}(t)$ or $\text{length}(s) = \text{length}(t)$ and s is lexicographically smaller than t .

Fix some $t \in \omega^{<\omega}$. Assume that we already decided which $s \triangleleft t$ will be elements of $T_{\max}$ as that we already defined the set of successors of all these s as well as the measure of their subsets. We have to define $\text{succ}(t)$ and the measure on it.

Let m_t be the number of nodes $s \triangleleft t$ already defined (including the successors of s for the already defined for $s \triangleleft t$). Set $M_t := (2m_t)^{m_t}$. Then we define $\text{succ}(t)$ to be of size $M_t^{m_t}$.¹⁵ For $A \subseteq \text{succ}(t)$ we set $\mu(A) := \log_{M_t}(|A|/M_t + 1)$.

¹⁴ $\mu(A \cup B) \leq \mu(A) + \mu(B)$.

¹⁵We can e.g. set $\text{succ}(t) := \{t \smallfrown k : 0 \leq k < M_t\}$.

Then $0 \leq \mu(A) < m_t$, $\mu(A) = 0$ iff $A = \emptyset$, and μ is strictly monotonous and sub-additive.¹⁶ If $A, B \subseteq \text{succ}(t)$ and $|B| \geq |A|/M_t$, then $\mu(B) > \mu(A) - 1$. If $|B| \leq m_t$ then $\mu(B) < 1/m_t$. If $\mu(\text{succ}(t)) > M$, then $m_t > M$.

Now fix an arbitrary $M \in \omega$. There is an N_0 such that $\mu(A) < 1/M$ for all s with $\text{length}(s) > N_0$ and all $A \subseteq \text{succ}(s)$ with $|A| < m_s$. (Just note that m_s strictly increases with $\text{length}(s)$.) Let N be larger than $M + 1$ and N_0 .

So assume that $s \perp t \in T_{\max}$, $\text{length}(s) > N \geq N_0$, $A \subseteq \text{succ}(t)$, $\mu(A) > N \geq M + 1$ (in particular $m_t > M$), and $f : A \rightarrow T_{\max}^{[s]}$.

Set $X := \{s' \geq s : s' \triangleleft t, \text{length}(s') \geq N\}$. Enumerate X as $\{s_0, \dots, s_{l-1}\}$ (for some $l \geq 0$). Set $A_0 := A$. Assume that A_n is already defined, and define

$$S_n := \{s' \in T_{\max} : (\exists t' \in A_n) s' \leq f(t')\}.$$

If $n > 0$ assume that $|\text{succ}_{S_n}(s_{n-1})| \leq 1$ and that $|A_n| > |A_{n-1}|/(2m_t)$.

Then we define A_{n+1} as follows: Since $s_n \in X$, $|\text{succ}(s_n)| < m_t$. By a simple pigeon-hole argument, there is an $A_{n+1} \subseteq A_n$ such that $|A_{n+1}| > |A_n|/(2m_t)$ and $|\text{succ}_{S_{n+1}}(s_n)| \leq 1$. So in the end we get a $B := A_l$ with cardinality at least $|A|/(2m_t)^{m_t} = |A|/M_t$, i.e. $\mu(B) > \mu(A) - 1 \geq M$. Also, $|\text{succ}_{S_l}(s')| \leq 1$ for every $s' \in X$, so $\mu_{S_l}(s') \leq 1/M$ (since $\text{length}(s')$ was sufficiently large).

We claim that B is as required. We have to show that S_l is $(N, 1/M)$ -meager. Pick an $s' \in S_l$ of length $\geq N$. We already dealt with the case $s' \in X$. Otherwise $s' \triangleright t$ (note that $s' \neq t$ since $s \perp t$). In this case $|\text{succ}_{S_l}(s')| \leq |\text{succ}_{T_{\max}}(t)| \leq m_{s'}$. So $\mu(\text{succ}_{S_l}(s')) \leq 1/M$, since $\text{length}(s') > N_0$. $\square$

Lemma 6.6. *If Q is strongly non-homogeneous, then P forces the following:*

If $r \in \lim(T_{\max}) \setminus \{\eta_i : i \in I\}$ then there is a $T \in V$ such that $r \in \lim(T)$ and T is $(1, 1)$ -meager.

If additionally the assumptions of Lemma 5.1 hold, then there are only $\aleph_1$ many $T \in V$, and $\aleph_1 < 2^{\aleph_0 V_{\omega^2}}$. This implies that the set $\{\eta_i : i \in I\}$ is of measure 1:

If $r \in \lim(T_{\max}) \setminus \{\eta_i : i \in I\}$, then $r \in \bigcup_{T \in V \text{ meager}} \lim(T) \in \mathbb{I}^c$.

Proof. Fix a P -name $\dot{r}$ for a real and a $p \in P$ such that $p \Vdash \dot{r} \notin \{\eta_i : i \in I\}$. We will show that there is a $p_\omega \leq p$ and a $(1, 1)$ -meager tree T such that $p_\omega \Vdash \dot{r} \in \lim(T)$.

We will by induction construct $p_n \in P$, approximations $g_n, k_n \in \omega$ and $i_n \in u_n = \text{dom}(g_n)$ such that

- (1) $p_0 = p$, $p_{n+1} \leq_{g_n} p_n$, g_{n+1} is purely stronger than g_n .
- (2) g_n is n -dense at i_n .
- (3) the sequence $(i_n)_{n \in \omega}$ covers $\bigcup \text{dom}(p_n)$ infinitely often.
- (4) $k_n \rightarrow \max(n+1, |\text{Pos}(g_n)|)$.
- (5) For each $\bar{a} \in \text{Pos}(g_n)$, $p_n^{[\bar{a}]}$ forces a value to $r \restriction k_n$. We call this value $r_n^{\bar{a}}$.
- (6) The tree $\{r_n^{\bar{a}} : \bar{a} \in \text{Pos}(g_n)\} \subseteq T_{\max} \restriction k_n$ is $(k_{n-1}, 1)$ -meager.


(1)–(3) allow us to fuse the $(p_n)_{n \in \omega}$ into a $p_\omega \leq p$ (cf. 3.11), and (5),(6) imply that the tree of all initial segments of r compatible with p_ω is meager.

So assume we have found p_n, g_n and k_n .

- Set $p := p_n$, $g := g_n$, $M := |\text{Pos}(g_n)|$ and $N := k_n$.
- Choose the position $i_{n+1} \in \text{dom}(p_n)$ according to some simple bookkeeping. This takes care of (3). Set $j := i_{n+1}$.
- Find a $q_1 \leq_g p_n$, $m > N$ such that $q_1 \Vdash (\eta_j \restriction m \neq \dot{r} \restriction m)$ and for all $\bar{a} \in \text{Pos}(g)$ $q_1^{[\bar{a}]}$ determines $\eta_j \restriction m$ and $\dot{r} \restriction m$:

First we apply pure decision 3.20(6) to get a $q' \leq_g p$ such that for all $\bar{a} \in \text{Pos}(g)$ there is an $m^{\bar{a}} > N$ and $\eta^* \neq r^*$ such that $q'^{[\bar{a}]} \Vdash (r^* \restriction m^{\bar{a}} = \eta^* \restriction m^{\bar{a}})$.

¹⁶Since the function $g(x) := \log_{M_t}(x+1)$ is concave and satisfies $g(0) = 0$.



F 7. h_2 (bold) is a subapprox. of h_1 and still purely stronger than g .

Then we apply pure decision again to get $q_1 \leq_g q'$ determining r and η_j up to $\max\{m^{\bar{a}} : \bar{a} \in \text{Pos}(g)\}$.

- Pick a q_1 -approximation h_1 which is N -dense at j and (purely) stronger than g .
- Pick a k_{n+1} such that $k_{n+1} \rightarrow \max(n+2, |\text{Pos}(h_1)|)$.
- Pick a $q_2 \leq_{h_1} q_1$ such that $q_2^{\lfloor \bar{a} \rfloor}$ determines η_j and r up to k_{n+1} for all $\bar{a} \in \text{Pos}(h_1)$.

So $q_2 \leq_g p$, h_1 approximates q_2 and witnesses N -density (at j). However, the tree of possible values for r could be very thick in the levels between k_n and k_{n+1} . We will thin out the approximation h_1 so that we still have $(n+1)$ -density, and the tree of possible values for r gets thin. (Recall that N is much larger than $n+1$, in particular $N \rightarrow n+1$.) We do this in two steps:

- Find a sub-approximation h_2 of h_1 that is still purely stronger than g and has only as many splittings as g , apart from one additional split (for each possibility) that witnesses N -density at j (see Figure 7).

To be more exact, we construct h_2 the following way: Given $\bar{a} \in \text{Pos}_{< j}(h_2)$. We have to define $h_2(i)(\bar{a})$. If $i \neq j$, pick for each $t \in g(i)(\bar{a} \upharpoonright g)$ exactly one successor $s \in h_1(i)(\bar{a})$. So h_2 makes the branches of g longer, but does not add any splittings. At j , we have the front $F := g(j)(\bar{a} \upharpoonright g)$ and the purely stronger N -dense front $F' := h_1(j)(\bar{a})$. Recall that $T := T_{\text{cldn}}^{F'} = \{s : s \leq F'\}$ is the finite tree corresponding to the front F' . We continue each $t \in F$ in T uniquely (without splits) until we reach a node with many (i.e. N -dense) splittings. We call this node “splitting node”. We take all the immediate successors of the splitting node and continue them uniquely in T until we reach a leaf of T , i.e. an element of F' . This process leads to a subset F'' of F' . Set $h_2(j)(\bar{a}) := F''$.

So we get: There are $|\text{Pos}_{\leq j}(g)| \leq M$ many pairs $(\bar{a}, t)$, where $\bar{a} \in \text{Pos}_{< j}(h_2)$ and t is a splitting node. Also, for $b \in \text{Pos}_{\leq j}(h_2)$, there are at most M continuations of b to some $b' \in \text{Pos}(h_2)$. Such a $b \in \text{Pos}_{\leq j}(h_2)$ corresponds to a pair $(\bar{a}, t)$ as above together with a choice of an (immediate) successor of t .

- Now we are ready to apply the Ramsey property. First fix a $\bar{a} \in \text{Pos}_{< j}(h_2)$ and a splitting node t . (There are at most M many such pairs.)

This pair corresponds to a unique $b \in \text{Pos}_{\leq j}(g)$. There are at most M many continuations of b to some $\bar{c} \in \text{Pos}(g)$. Fix an enumeration $\bar{c}_1 \dots \bar{c}_M$ of these possible continuations. Each $\bar{c}_l$ forces a value to $r \upharpoonright k_n$, call this value r^l .

Back to h_2 . Set $A := \text{succ}(t)$ in the tree $T_{\text{cldn}}^{h_2(j)(\bar{a})}$ (or equivalently $T_{\text{cldn}}^{h_1(j)(\bar{a})}$). So $\mu(A) \geq N$. For every $s \in A$ there is a unique $s' \geq s$ such that $\bar{a} \cup \{(j, s')\} \in \text{Pos}_{\leq j}(h_2)$, and for every $s \in A$, $l \in M$ there is a unique $b \in \text{Pos}(h_2)$ continuing $\bar{c}_l \in \text{Pos}(g)$ and $\bar{a} \cup \{(j, s')\}$. Each such b decides r up to k_{n+1} . We call this value $r^{s,l}$. So $r^{s,l} \upharpoonright k_n = r^l$. Also $r^l \perp t$ since t is a possible initial segment of η_j .

For every $l \in M$ we define a function $f_l : A \rightarrow T_{\max}^{[r^l]} \upharpoonright k_{n+1}$ by mapping s to $r^{s,l}$. So we can apply the Ramsey property and get a $B \subseteq A$ such that $\mu(B) > M \geq n+1$, and the tree of possibilities for $\dot{r}$ induced by $\bar{a}, B$ is $1/M$ -meager. We repeat that for all pairs $(\bar{a}, t)$ where $\bar{a} \in \text{Pos}_{< j}(\mathfrak{h}_2)$ and t is a splitting node, and get a subapproximation $\mathfrak{g}_{n+1}$ of $\mathfrak{h}_2$ such that the tree of possibilities for $\dot{r}$ induced by $\mathfrak{g}_{n+1}$ is $(k_n, 1)$ -meager (here we again use the sub-additivity of μ).

This results in a sub-approximation $\mathfrak{g}_{n+1}$ of $\mathfrak{h}_2$ (and therefore $\mathfrak{h}_1$) which is still purely stronger than $\mathfrak{g} = \mathfrak{g}_n$. Since $\mathfrak{g}_{n+1}$ is a sub-approximation of $\mathfrak{h}_1$, $|\text{Pos}(\mathfrak{g}_{n+1})| \leq |\text{Pos}(\mathfrak{h}_1)|$, and therefore $k_{n+1}, \mathfrak{g}_{n+1}$ satisfy (4). $\square$

Note that we did not use the j_α or automorphisms of I , the proof works for all I_0 . $I = \{i\}$ in particular implies: If $Q \Vdash \dot{r} \neq \dot{\eta}_i$, then for all $T \in Q$ there is a $S \leq_Q T$ such that $r \notin \lim(S)$, i.e.

Corollary 6.7. *If Q is strongly non-homogeneous then Q forces that $\dot{\eta}$ is the only Q -generic real over V in $V[G_Q]$.*

Remark. A similar forcing Q^{JeSh} (finitely splitting, rapidly increasing number of successors) was used in [4] to construct a complete Boolean algebra without proper atomless complete subalgebra. Q^{JeSh} can also be written as lim-sup forcing. However, the difference is that the norm in Q^{JeSh} is “discrete” (as e.g. Sacks): either s has a minimum number of successors, then the norm is large, or the norm is < 1 . Such a norm cannot satisfy a Ramsey property as the one above. For Q^{JeSh} we can only prove Corollary 6.7, but not Lemma 5.9.

We have already mentioned another corollary:

Corollary 6.8. *If Q is strongly non-homogeneous, then P forces that $\{\dot{\eta}_i : i \in I\}$ is of measure 1.*

This, together with 5.7 and 5.9 proves Theorem 4.7.

Remark. There are various ways to extend the constructions in this paper. As already mentioned, we could use non-total orders I or allow Q_i to be a $P_{< i}$ -name. A more difficult change would be to use lim-inf trees instead of lim-sup trees. In this case we need additional assumptions such as bigness and halving. This could allow us to apply saccharinity to a ccc ideal $\mathbb{I}$, i.e. to force (without inaccessible or amalgamation) measurability of all definable sets.

R

- [1] Jörg Brendle. Mad families and iteration theory. In Yi Zhang, editor, *Logic and algebra*, volume 302 of *Contemporary Mathematics*, pages 1–31. Amer. Math. Soc., Providence, RI, USA, 2002.
- [2] Marcia Groszek. ω_1^* as an initial segment of the c -degrees. *J. Symbolic Logic*, 59(3):956–976, 1994.
- [3] Marcia Groszek and Thomas Jech. Generalized iteration of forcing. *Transactions of the American Mathematical Society*, 324, No. 1.:1–26, 1991.
- [4] Thomas Jech and Saharon Shelah. A complete Boolean algebra that has no proper atomless complete subalgebra. *Journal of Algebra*, 182, 1996. math.LO/9501206.
- [5] Vladimir Kanovei. On non-wellfounded iterations of the perfect set forcing. *The Journal of Symbolic Logic*, 64, No. 2.:551–574, 1999.
- [6] Andrzej Rosłanowski and Saharon Shelah. Norms on possibilities I: forcing with trees and creatures. *Memiors of the American Mathematical Society*, 141(671):xii + 167, 1999. math.LO/9807172.
- [7] Andrzej Rosłanowski and Saharon Shelah. Sweet & Sour and other flavours of ccc forcing notions. *Archive for Mathematical Logic*, 43:583–663, 2004. math.LO/9909115.
- [8] Saharon Shelah. Two cardinal invariants of the continuum ($\mathfrak{d} < \mathfrak{a}$) and FS linearly ordered iterated forcing. *Acta Mathematica*, accepted. Also known under the title “Are $\mathfrak{a}$ and $\mathfrak{d}$ your cup of tea?”. math.LO/0012170.
- [9] Saharon Shelah. Can you take Solovay’s inaccessible away? *Israel Journal of Mathematics*, 48:1–47, 1984.
- [10] Robert M. Solovay. A model of set theory in which every set of reals is Lebesgue measurable. *Annals of Math.*, 92:1–56, 1970.

J K , I " D M G , T U " W , 1050 W ,
A

E-mail address: kellner@fsmat.at

URL: <http://www.logic.univie.ac.at/~kellner>

S S , E I M , E J. S C , G R , T H
U J , J , 91904, I , D M , R U , N
B , NJ 08854, USA

E-mail address: shelah@math.huji.ac.il

URL: <http://www.math.rutgers.edu/~shelah>